

## ***Amphizonella violacea* Greeff, 1866**

**Most likely ID:** n.a.


**Synonym:** n.a.

**Sampling location:** Moss

**Phylogenetic tree:** [Amphizonella violacea](#)

### **Diagnosis:**

- body globular, very slow movement
- pseudopodia hyaline and clear
- diameter 125–250  $\mu\text{m}$
- soft and flexible shell, thickness 3–23  $\mu\text{m}$
- pseudostome irregularly shaped
- cytoplasm filled with numerous violet vesicles
- one ellipsoid or globular nucleus, diameter up to 40  $\mu\text{m}$
- about 20 contractile vacuoles near surface



### *Amphizonella violacea*

So far, I have found *Amphizonella violacea* exclusively in mosses on roofs. After moistening the moss samples with collected rainwater, the first excysted specimens were found after 2 days. In 2006, Meisterfeld and Badewitz published a redescription of *Amphizonella violacea*. The authors also found their specimens in mosses on roofs. However, other authors such as Siemensma (s. link below) and Glück (2007) report that *Amphizonella violacea* can also be found in *Sphagnum* bogs.

At low magnifications, the rounded specimens are easy to overlook because they often appear black and look like lifeless detritus particles. Only at higher magnification and with slight coverslip pressure can the distinct violet coloration of *Amphizonella violacea* be recognized. This coloration is produced by large amounts of violet-stained vesicles with a diameter of 0.1–2  $\mu\text{m}$ , which are distributed throughout the cytoplasm (s. fig. 6).

The specimens of *Amphizonella violacea* moved very little and then only very

slowly. Most of the time, the body was rounded or knobby. The rounded specimens had a diameter of 160–230  $\mu\text{m}$ . The specimens phagocytized large amounts of detritus, algae, and also mineral grains. Everything was phagocytized completely non-specifically.

In all specimens of my population, I could only identify one nucleus. This was ellipsoid or round in shape with a diameter of 35–40  $\mu\text{m}$ . Numerous, flake-shaped nucleoli were visible in the periphery (s. fig. 3 a-b). This corresponds with the observations of Meisterfeld & Badewitz (2006), Glück (2007), and Siemensma (s. link below). Only Penard (1902) reported finding specimens with 2 or 3 nuclei. In slightly compressed specimens, numerous contractile vacuoles can be seen, all located just beneath the cell membrane.

The shell of *Amphizonella violacea* consists of a soft and flexible, chitinous layer. In some specimens, I was able to observe layering. The shell was either colorless or distinctly yellow-brown (s. figs 5 and 7). In some cases, I also observed how the shell clearly detached from the cell body, creating blister-like elevations (s. fig. 7). This detachment of the shell may be caused by the movement of the cell body, which the shell cannot adapt to. However, this phenomenon illustrates that the shell lies loosely against the plasma body. The thickness of the shell in *Amphizonella violacea* seems to be subject to some variability. I measured a range of 1–5  $\mu\text{m}$ . Meisterfeld and Badewitz report a thickness of up to 25  $\mu\text{m}$ . The shell can be covered by a gelatinous layer, in which bacteria are often found. However, I could not detect this layer in all specimens, which corresponds with the observations of Siemensma (s. link below).

*Amphizonella violacea* is difficult to distinguish from the very similar, also violet-colored species [Zonomyxa violacea](#). Meisterfeld and Badewitz (2006) cite the following essential differences:

- *Amphiozella violacea* has only one nucleus, while [Zonomyxa violacea](#) has 4–32
- a gelatinous layer over the shell is absent in [Zonomyxa violacea](#)
- [Zonomyxa violacea](#) is said to have a thinner shell than *Amphizonella violacea*
- the habitat of [Zonomyxa violacea](#) is *Sphagnum* bogs, while *Amphizonella violacea* lives in xerophilous mosses (on roofs, walls, or trees) with strongly fluctuating water content

However, in my opinion, some of these characteristics are very unspecific. The habitats overlap, as *Amphizonella violacea* has also been found in *Sphagnum* bogs. The thickness of the shell is very variable in both species, and here too the measurements overlap. Finally, the gelatinous sheath can also be absent in *Amphizonella violacea*. The number of nuclei remains the most reliable characteristic. *Amphizonella violacea* always seems to be uninucleate.

More images and information on *Amphizonella violacea*: [Ferry Siemensma-Microworld-Amphizonella violacea](#)

Amphizonella violacea  
Obj. 60 X




© Martin Kreutz

**Fig. 1:** *Amphizonella violacea*. D = 158  $\mu$ m. A rounded, resting specimen. Obj. 60 X.

Amphizonella violacea  
Obj. 60 X




**Fig. 2:** *Amphizonella violacea*. D = 158  $\mu$ m. In a slightly squashed specimen the single, globular nucleus (Nu) is visible. SH = shell. Obj. 60 X.



**Fig. 3 a-b:** *Amphizonella violacea*. Two focal planes of the nucleus (Nu) with numerous peripher nucleoli (Nuc). The nucleus has a diameter of 40  $\mu\text{m}$ . Obj. 100 X.

Amphizonella violacea  
Obj. 40 X




**Fig. 4:** *Amphizonella violacea*. D = 220  $\mu$ m. Ventral view on the irregularly shaped pseudostome (PS). Obj. 40 X.

Amphizonella violacea  
Obj. 100 X



**Fig. 5:** *Amphizonella violacea*. The chitinoid shell (SH) of this specimen has a thickness of about 3  $\mu\text{m}$ . A gelatinous sheath is absent. CV = contractile vacuoles. Obj. 100 X.



**Fig. 6:** *Amphizonella violacea*. The violet vesicles scattered in the cytoplasm have a diameter of 0.5 to 1.5  $\mu\text{m}$ . Obj. 100 X.

Amphizonella violacea  
Obj. 60 X



**Fig. 7:** *Amphizonella violacea*. In some specimens the shell was colored yellowish-brown and detached from the cell surface in a blister-shape (arrows). CV = contractile vacuoles. Obj. 60 X.