

***Aphanothece stagnina***  
**(Sprengel) A.Braun, 1863**

**Most likely ID:** n.a.


**Synonym:** *Coccochloris stagnina*

**Sampling location:** [Simmelried](#), [Sima Moor \(Austria\)](#)

**Phylogenetic tree:** [Aphanothece stagnina](#)

**Diagnosis:**

- colonies in a common mucilaginous sheath
- young colonies spherical, with distinct margin
- older colonies irregular, up to 4 cm in diameter
- cells oval or cylindrical, rounded ends
- length (of cells) 3.8–9 µm, width 3–5 µm
- no individual mucilaginous envelope of cells
- color bluish-green
- cells loosely dispersed in colony




### Aphanothece stagnina

The cyanobacterium *Aphanothece stagnina* forms round or irregularly shaped gelatinous colonies, which usually lie on the bottom mud. So far I have found this species in the [Simmelried](#) and in the [Sima Moor](#) in Austria.


The cells of *Aphanothece stagnina* are oval or cylindrical and between 4–9  $\mu\text{m}$  long. Rarely, cells up to 11  $\mu\text{m}$  are found, whereby one must ensure that no dividers are measured. In my population the cells were mostly 7–8  $\mu\text{m}$  long and about 5  $\mu\text{m}$  wide. The largest colonies I found had a diameter of about 500  $\mu\text{m}$  and were irregularly shaped. I have never found macroscopic colonies up to several centimeters in diameter.

The differentiation from the very similar species *Aphanothece microscopica* is insufficiently defined. The cell shape and also the size of the cells are practically identical. Komarek & Anagnostidis (1999) cite the size of the colonies as a distinguishing feature. In *Aphanothece stagnina* it is said to reach 4–6 cm, whereas the colonies of *Aphanothece microscopica* are said to reach a maximum of 2 mm. However, Huber-Pestalozzi (1938) mentions the width of the cells as a distinguishing feature. The cells of *Aphanothece microscopica* are said to be 2–3  $\mu\text{m}$  wide, while those of *Aphanothece stagnina* are said to be 3–6.5  $\mu\text{m}$  wide. In addition, according to Huber-Pestalozzi, the cells of *Aphanothece microscopica* are said to have individual mucous membranes, while the cells of *Aphanothece stagnina* are embedded in a homogeneous mucus. In my opinion, the size of the colonies is not a good determinant because young colonies must also be taken into account. I therefore agree with the

distinguishing features of Huber-Pestalozzi. The cells of my population had a width of mostly 5  $\mu\text{m}$  and I could not detect individual mucus sheaths around the cells. It must therefore be *Aphanothece stagnina*.




**Fig. 1 a-b:** *Aphanothece stagnina*. D = 125  $\mu\text{m}$  (of colony). Two focal planes of a young, spherical colony embedded in detritus. MC = margin of the colony. Obj. 60 X.




**Fig. 2:** *Aphanothece stagnina*. L = 7-8 µm (of cells). The cells of the colony as shown in fig. 1 a-b in detail. Obj. 100 X.






**Fig. 3:** *Aphanothece stagnina*. L = 7-8  $\mu\text{m}$  (of cells). The cells shown in fig. 2 in brightfield illumination. Note the fine, dark granules in the cytoplasm of the cells. Obj. 100 X.



**Fig. 4 a-b:** *Aphanothece stagnina*. L = 5–9.5  $\mu\text{m}$  (of cells). Two focal planes of a second, small colony in DIC. MC = margin of the colony. Obj. 100 X.





**Fig. 5 a-b:** *Aphanothece stagnina*. L = 5-9.5  $\mu\text{m}$  (of cells). The colony as shown in fig. 4 a-b in brightfield illumination. Obj. 100 X