

Archerella flavum

(Archer, 1877) Loeblich & Tappan, 1961

Most likely ID: n.a.


Synonym: *Amphitrema flavum*

Sampling location: [Sima Moor \(Austria\)](#)

Phylogenetic tree: [Archerella flavum](#)

Diagnosis:

- shell ovoid to elliptical, compressed, brownish transparent
- shell with two elliptical apertures at opposite ends
- length 45–77 µm
- several contractile vacuoles
- nucleus spherical with central nucleolus
- filopodia thin, straight, sometimes branched
- symbiotic algae present




Archerella flavum

I found *Archerella flavum* in a small puddle in the Sima Moor (Austria) in June 2024. This filose testate amoeba is easy to identify by its conspicuous shell. It is usually brownish in color, transparent, without attached xenosomes and has two opposite apertures. The protoplast can be seen clearly through the transparent shell. It is attached to the inner wall of the shell with strands of cytoplasm. The thin, mostly straight filopodia are only extended by undisturbed specimens (s. figs. 1 a and 2 a).


The protoplast of *Archerella flavum* is usually clearly green in color because this testate amoeba has symbiotic algae. These algae have never been described in detail in the literature available to me. According to my observations, the algae cells are spherical with a diameter of 4.0-4.5 μm . They seem to be of the *Chlorella* type. They have a cup-shaped chloroplast, each with a pyrenoid. I could also see that each algae cell has a separate nucleus.

Further images and information on *Archerella flavum*: [Ferry Siemensma-Microworld-Archerella flavum](#)

Archerella flavum
Obj. 100 X



a



b

Fig. 1 a-b: *Archerella flavum*. L = 50 μ m. Two focal planes of an unsquashed specimen. Note the filopodia (Fi), which are extended from the two apertures of the shell. CV = contractile vacuoles, Nu = nucleus, SA = symbiotic algae. Obj. 100 X.

Archerella flavum
Obj. 100 X



Fig. 2 a-b: *Archerella flavum*. L = 53 μ m. Two focal planes of a second specimen with extended filopodia. Obj. 100 X.




Fig. 3: *Archerella flavum*. L = 53 µm. The slightly squashed specimen as shown in fig. 2 a-b to show the symbiotic algae (SA) in detail. The algae seem to be of the *Chlorella* type with a cup-shaped chloroplast with a single pyrenoid (PY). Obj. 100 X.