## Blepharisma coeruleum (Gajevskaja, 1927)

Most likely ID: n.a.

Synonym: n.a.

**Sampling location:** <u>Simmelried</u>

Phylogenetic tree: **Blepharisma coeruleum** 

## **Diagnosis:**

- length 120 145 µm
- cell teardrop-shaped, anterior end flattened
- CV terminal
- macronucleus spherical to oval with 3-6 adjacent micronuclei
- Granula bluish or green-bluish
- contains symbiotic algae
- adorale zone reaches third of body



Fig. 1: Blepharisma coeruleum. L =  $105 \mu$ m, ventral view of a freely swimming specimen.



Fig. 2: Blepharisma coeruleum. L =  $105 \mu$ m, ventral view with focus on the spherical macronucleus. Obj. 100 X.



**Fig. 3:** Blepharisma coeruleum. L =  $105 \mu$ m, ventral view with focus on the rows of pink granula. Obj. 100 X.

The anteriorly pointed shape and the pink coloration are very typical of *Blepharisma*. The length of the specimens observed varied between  $100 - 120 \mu m$ , which is quite small for *Blepharisma coeruleum*. In fact, the specimens shown in Fig. 1 are very similar to

*Blepharisma lateritium,* which is supposed to grow to a similar size:



Blepharisma lateritium

However, all of the specimens I have observed (about 40) contained symbiotic algae that filled the posterior half of the body. This is not described for *Blepharisma lateritium*. According to the original description of *Blepharisma coeruleum* by Gajevskaja this species should have a distinct bluish or blue-green coloration. This was not the case in my specimens, but the description of Gajevskaja was adopted by later authors (Kahl, Foissner) without having examined the species itself *in vivo*. Kahl pointed out the similarity between *Blepharisma coeruleum* and *Blepharisma lateritium* and also considers it a variant of *Blepharisma lateritium*. Kahl further writes (I assume according to Gajevskaja's description): "apparently eats only one particular alga (*Tetraspora*)". In fact, the algae in my specimens were all of the same species and look very similar to *Tetraspora* (s. Fig. 4). However, the algae are certainly not ingested, but symbiotic algae. I conclude this on the basis of the following observations:

– all observed specimens had about the same number of algae in the plasma, always in the posterior end

- all algae are of the same species

- the clearly visible food vacuoles of the specimens exclusively contained bacteria (s. Fig. 5)

- the algae are not sequestered in separate vacuoles



**Fig. 4:** Blepharisma coeruleum. The symbiotic algae with a diameter of 4-6  $\mu$ m are not of the Chlorella type but similar to Tetraspora. Obj. 100 X



**Fig. 5**: *Blepharisma coeruleum*. Squashed specimen. FV = food vacuoles, SA = symbiotic algae. Obj. 100 X.

The presence of symbiotic algae confirms the identification as *Blepharisma coeruleum*. However, there is some uncertainty regarding the state of the nuclear apparatus. Gajevskaja does not describe a number of micronuclei. For the similar species *Blepharisma lateritium*  there are different data. Kahl indicates the presence of only one micronucleus in this species, while Foissner describes "many small" micronuclei. This may mean that the number of micronuclei in *Blepharisma lateritium* is variable. In my population specimens with 3 – 6 micronuclei were present:



**Fig. 6:** *Blepharisma coeruleum*. The macronucleus with 4 attached micronuclei. Mi = micronuclei, Ma, = macronucleus. Obj. 100 X.

A further uncertainity is the color of the granula. Gajevskaja described a blue-green coloration. Therefore I examined the ciliate again in brightfield, as Gajevskaja probably did:





The image in Fig. 7 indicates clearly a pink color. However, if the aperture is closed a bit, they appear almost black. It is difficult to say under what conditions and with what kind of optical equipment Gajevskaja investigated the color of the granules at that time, but my specimens were definitely not blue or blue-green. It is known, however, that the dye in the granules (Blepharismin) can change color when exposed to light and oxygen. These facts demonstrate the difficulties in comparing present day observations with the descriptions and drawings of earlier authors. In my opinion, *Blepharisma coeruleum* with a pink coloration is present here, or *Blepharisma lateritium* with symbiotic algae. Because

*Blepharisma lateritium* was found and described several times (in contrast to *Blepharisma coeruleum*) and symbiotic algae were never observed, I stick to the classification *Blepharisma coeruleum*.