

## ***Brachonella pulchra***

(Kahl, 1927) Bourland et al. 2018

**Most likely ID:** n.a.


**Synonym:** *Metopus pulcher*

**Sampling location:** [Simmelried](#)

**Phylogenetic tree:** [Brachonella pulchra](#)

### **Diagnosis:**

- body broadly ellipsoidal or obovoidal
- length 80–110 µm
- adoral zone running almost a full turn around longitudinal axis
- on ventral side adoral zone runs diagonally before it bends to continue in parallel to longitudinal axis
- mouth opening in posterior fourth of body
- macronucleus globular
- fringe of inconspicuous extrusomes beneath pellicle
- contractile vacuole terminal
- posterior end broadly rounded
- slightly elongated caudal cilia




1, 2 = after Kahl  
3, 4 = after Bourland et al.

### **Brachonella pulchra**

I have found *Brachonella pulchra* so far exclusively in the uppermost mud layer in [Simmelried](#). The species is not very common. In the last 20 years I could observe only a few specimens. I recognize *Brachonella pulchra* by the very posteriorly located mouth opening


(s. fig. 1 b) and the course of the adoral zone, which runs almost around the entire body and suddenly turns posteriorly on the ventral side, almost in parallel to the longitudinal axis. At the broadly rounded posterior end a few, slightly elongated caudal cilia arise, which can be easily overlooked. As already described by Kahl, the species is quite variable. So I could observe a smaller form in December 1998, which was only 84  $\mu\text{m}$  long (s. fig. 2 a-d).



**Fig. 1 a-f:** *Brachonella pulchra*. L = 108  $\mu\text{m}$ . A freely swimming specimen from ventral (a, b), from right (c, d) and from dorsal (e, f). AZM = adoral zone of membranelles, CC = caudal cilia, CV = contractile vacuole, Ma = macronucleus, MO = mouth opening, PC = perizonal cilia. Obj. 60 X



*Brachonella pulchra*  
Obj. 40 X



**Fig. 2 a-d:** *Brachonella pulchra*. L = 84  $\mu\text{m}$ . A second, freely swimming specimen from ventral. Obj. 40 X