Colpoda cucullus

(Müller, 1773) Gmelin, 1790

Most likely ID: n.a.

Synonym: n.a.

Sampling location: Moss

Phylogenetic tree: Colpoda cucullus

Diagnosis:

- body broadly reniform, laterally slightly flattened
- semicircular indented at the mouth opening almost in mid-body
- length 40–120 μm , usually 60–80 μm
- macronucleus globular to slighty ellipsoid, condensed chromatin sometimes starshaped
- lenticular micronucleus adjacent to macronucleus, about 3 x 2 μm , sometimes 2–4 micronuclei present
- mouth opening with a right and left field of polykineties
- extrusomes 1–2 μm long, comma-shaped, moderately refractive
- ciliary rows consiting of paired cilia
- contractile vacuole terminal
- no caudal cilia



Colpoda cucullus

I found my population of *Colpoda cucullus* in moss samples that came mainly from trees and walls. If the moos is poured over with rainwater in Petri dishes, a large population forms after a few days. *Colpoda cucullus* is actually the typical ciliate in hay infusions, where it practically always occurs. The habitats and reproduction strategies of *Colpoda cucullus* were studied in detail by Mueller & Muller (1970). According to this study, cysts of *Colpoda cucullus* are found on practically all green plants. They are activated by dew or rainwater. The life cycle of *Colpoda cucullus* is short enough for a complete reproduction cycle before the leaf surfaces dry again. The cysts are transferred to other plants by pollen-collecting insects.

Colpoda cucullus is easily confused with *Colpoda lucida* in terms of size and shape. But *Colpoda lucida* has a clear, light-colored fringe of densely packed extrusomes, which makes it easy to distinguish. In my populations, the specimens of *Colpoda cucullus* were about 50-70 µm long, which is exactly in the lower range of the typical range given by Foissner. The macronucleus has no nucleoli, but is criss-crossed by chromatin islands, which can be clearly seen in squashed specimens (s. fig. 4). To the right and left of the mouth opening are two dense fields of cilia called polykinetids. In ventral view, these two polykinetids are clearly visible (s. fig. 3 a-b).



Fig. 1 a-d: Colpoda cucullus. L = 62 μ m. Different focal planes of a freely swimming specimen from right. Obj. 60 X.



Fig. 2 a-d: Colpoda cucullus. $L = 70 \mu m$. Different focal planes of a slightly squashed specimen from right. Note the cloudy chromatin in the macronucleus (Ma). CV = contractile vacuole, Mo = mouth opening. Obj. 100 X.



Fig. 3 a-b: Colpoda cucullus. $L = 51 \mu m$. A squashed specimen from ventral. Note the right oral polykinetid (ROP) and the left oral polykinetid (LOP). Obj. 100 X



Fig. 4: Colpoda cucullus. A strongly squashed specimen. CV = contractile vacuole, FV = food vacuoles, Ma = macronucleus with chromatin, Mi = micronucleus. Obj. 100 X



Fig. 5: Colpoda cucullus. The comma-shaped extrusomes (EX) in a strongly squashed specimen. Obj. 100 X