Colpoda steinii Maupas, 1883

Most likely ID: n.a.


Synonym: Colpoda steini

Sampling location: Moss

Phylogenetic tree: Colpoda steinii

Diagnosis:

- body reniform, dorsal side convex, ventral side almost straight
- length 10-60 μm, usually 20-40 μm
- semicircularly indented at the mouth opening
- apical keel with 5-7 ribs
- macronucleus ellipsoid with one central nucleolus
- mouth opening with a right and left field of polykineties
- cilia of the left polykinetid are long, forming a "beard"
- 10-13 ciliary rows of paired cilia
- contractile vacuole terminal
- two caudal cilia (hard to see)



Colpoda steinii

I find *Colpoda steinii* in almost all moss samples that I soak with rainwater in Petri dishes. After a few days, populations of varying sizes occur. However, Colpoda steinii can also be found in hay infusions, soil samples and even in plankton.

Colpoda steinii can easily be confused with other Colpoda species, such as Colpoda ecaudata or Colpoda aspera. Colpoda ecauda, however, has several nucleoli in the macronucleus and not just a central one. Colpoda aspera is smaller and has a pointed apical end. Both species also have no caudal cilia, but Colpoda steinii has.

The two caudal cilia of *Colpoda steinii* are hard to see at high magnifications because they are constantly moving and are rarely in focus. It is therefore advisable to examine specimens at first at lower magnifications. The characteristic "beard", which originates in the middle of the ventral side, is also striking. These are the long cilia of the left oral cilia. This dense field of cilia (polykinetid) is difficult to recognize under the light microscope. However, it is possible to distinguish between the left and right cilia of the mouth opening when focusing through the specimen.




Fig. 1 a-b: Colpoda steinii. L = 28 $\mu m.$ A freely swimming specimen from left. Note the two caudal cilia (CC). Obj. 40 X.




Fig. 2 a-d: Colpoda steinii. L = 33 μm. A second freely swimming specimen from right. Note the "beard" (BE) of long cilia of the left oral polykinetid. CC = caudal cilia. Obj. 60 X.



Fig. 3 a-c: Colpoda steinii. L = 39 µm. A slightly squashed specimen from right. Note the right oral polykinetid (ROP) and the left oral polykinetid (LOP). In the center of the macronucleus (Ma) a spherical nucleolus is visible. AK = apical keel with ribs, CV = contractile vacuole, PC = paired cilia. Obj. 100 X.



Fig. 4: Colpoda steinii. $L=34~\mu m$. A slightly squashed specimen from left. LOP = left oral polykinetid, Nuc = nucleolus (in center of macronucleus). Obj. 100 X.