

***Desmidium grevillei***

**(Ralfs) De Bary, 1858**

**Most likely ID:** n.a.


**Synonym:** n.a.

**Sampling location:** [Schwemm Moor \(Austria\)](#)

**Phylogenetic tree:** [Desmidium grevillei](#)

**Diagnosis:**

- unbranched filaments of almost rectangular cells
- cells in apical view elliptical
- filaments are twisted, covered by mucilage
- cells 40–55 µm wide, 20–25 µm long
- cells slightly constricted in the middle
- basal angles of semi-cells conical
- each semi-cell with a ring of granules
- one chloroplast per semi-cell with one pyrenoid



after Lenzenweger


*Desmidium grevillei*

So far, I have only found *Desmidium grevillei* in the [Schwemm Moor](#) in Austria. The algae was very common in the samples.


The cells in the filaments of *Desmidium grevillei* consist of two semi-cells, like many other desmid algae (e.g. *Cosmarium* or *Staurastrum*). However, the isthmus is only very weakly pronounced and is only a slight tapering. Each semi-cell has conical basal angles. Here, the cell wall is also significantly thickened. Since the cells in the filament are always slightly offset from each other, these conical thickenings appear to run spirally around the filament (s. fig. 4). Each semi-cell also has a ring of fine pores, which are only visible under high

magnification (s. fig. 5).

In lateral view, the individual cells appear almost rectangular. They are connected to each other with their apices. If a cell is detached from the filament and rotated, it can be viewed in apical view (s. fig. 3 a-b). The shape is similar to that of a lemon. The conical ends are the basal angles of the semi-cells.




**Fig. 1 a-b:** *Desmidium grevillei*. L = 30-32  $\mu\text{m}$  (of cells). Two focal planes of the end of a filament in DIC. Obj. 40 X.




**Fig. 2 a-b:** *Desmidium grevillei*. L = 30-32  $\mu\text{m}$  (of cells). The same specimen as shown in fig. 1 a-b in brightfield illumination. Obj. 40 X.






**Fig. 3 a-b:** *Desmidium grevillei*. Two focal planes of a slightly squashed end of a filament with a separated cell (SC). The separate cell is visible in apical view with an elliptical shape

and conical shaped basal angles. Obj. 100 X.



**Fig. 4:** *Desmidium grevillei*. The cells in the filament are arranged with an offset to each other. The focal planes is on the conical shaped basal angles (BA) of the cells. Obj. 40 X.



**Fig. 5:** *Desmidium grevillei*. The cell wall of the semi-cells are covered with a ring of granules (GR). Obj. 100 X.