

Dexiotricha elliptica

(Kahl, 1931) Fan et al., 2014

Most likely ID: n.a.


Synonym: *Loxocephalus ellipticus*

Sampling location: [Ulmisried](#), [Purren pond](#)

Phylogenetic tree: [Dexiotricha elliptica](#)

Diagnosis:

- body long ellipsoidal with unciliated frontal plate
- length 40-55 µm
- oral apparatus small, subapical
- globular macronucleus below cell equator
- one micronucleus adjacent to macronucleus
- extrusomes spindle-shaped, about 4 µm long
- contractile vacuole in posterior fifth
- cytoplasm filled with refractile granules, not ring-shaped
- granules arranged in rows beneath pellicle, 8-9 rows per body side
- one long caudal cilium



Dexiotricha elliptica

Dexiotricha elliptica was first described by Kahl in 1931 as *Loxocephalus ellipticus*. In 2014, Fan et al. transferred the species to the genus *Dexiotricha* on the basis of morphological investigations and renamed it *Dexiotricha elliptica*.

I find *Dexiotricha elliptica* quite frequently in the mud layer of decomposed leaves in [Ulmisried](#) and [Purren pond](#). The ciliate is filled with granules, which are never ring-shaped in my specimens. Instead, the granules are mostly spherical or slightly asymmetrical in shape with a maximum diameter of 1.5 µm. Under the pellicle, the granules are arranged in longitudinal rows, as described by Kahl. IN my population the macronucleus was always located below the cell equator, and the contractile vacuole in the posterior fifth. I was also able to recognize the short canal between the contractile vacuole and the excretory porus on the ventral side, as drawn by Kahl (s. drawing 3 above).

Dexiotriche elliptica differs from the similar species [Loxocephalus moniligranulatus](#) and *Dexiotricha granulosa* mainly in the shape of the granules and the position of the contractile vacuole. In [Loxocephalus moniligranulatus](#) the contractile vacuole is located in the middle of

the body and in *Dexiotricha granulosa* the granules are ring-shaped and the contractile vacuole is also located in the middle of the body.




Fig. 1 a-d: *Dexiotricha elliptica*. L = 52 μ m. A freely swimming specimen. Note the short excretion canal (EC) of the contractile vacuole. CC = caudal cilium, CV = contractile vacuole, FP =frontal plate, Ma = macronucleus, MO = mouth opening. Obj. 60 X.




Fig. 2 a-c: *Dextotricha elliptica*. L = 52 μ m. The slightly squashed specimen as shown in fig. 1 a-d. Note the granula (GR) arranged in longitudinal rows beneath the pellice. CV = contractile vacuole, Ma = macronucleus. Obj. 100 X.




Fig. 3 a-d: *Dexiotricha elliptica*. L = 59 μm . Different focal planes of a second, slightly squashed specimen. CC = caudal cilium, CV = contractile vacuole, GR = granules, Ma = macronucleus, MO = mouth opening. Obj. 60 X.




Fig. 4 a-d: *Dexiotricha elliptica*. Different focal planes of a strongly squashed specimen. CC = caudal cilium, CV = contractile vacuole, GR = granules, Ma = macronucleus, Mi? = probably the micronucleus, MO = mouth opening. Obj. 100 X.




Fig. 5 a-b: *Dexiotricha elliptica*. A second strongly squashed specimen. Note the excretion pore (EP) of the contractile vacuole. CV = contractile vacuole, Ma = macronucleus, Mi = micronucleus, MO = mouth opening. Obj. 100 X.