

***Diaphoropodon mobile* Archer, 1869**

Most likely ID: n.a.


Synonym: n.a.

Sampling location: [Sima Moor \(Austria\)](#), [Schwemm \(Austria\)](#), [Simmelried](#)

Phylogenetic tree: [Diaphoropodon mobile](#)

Diagnosis:

- test ovoid or pear-shaped, membranous and flexible, brownish color
- test covered with organic and inorganic material
- hyaline, transparent rods adhere to the test, 8–10 µm long
- length 25–178 µm
- aperture of test broad, circular or oval
- nucleus in posterior end of the test
- one or two contractile vacuoles
- filipodia branching and anastomosing, arising in all directions



after Penard

Diaphoropodon mobile

I have found *Diaphoropodon mobile* so far only in the Sima Moor (Austria) in the Schwemm (Moor, Austria) and in the [Simmelried](#). However, in [Simmelried](#) I have not been able to detect the species in the last 7 years.

The most conspicuous feature of *Diaphoropodon mobile* are thin, hyaline rods with which the test is covered. The purpose of these rods is unknown. They are absent in empty tests of dead specimens. Thus, it is possible that these rods are formed by the amoeba and protrude through the test. The length of the rods is reported to be 8-10 μm . However, in the specimens I examined, the length varied considerably between 4-18 μm (s. figs. 1, 2 and 3 a-b). If the rods are really different in length or if they protrude through the test with different lengths due to different thicknesses of the test cannot be determined.

In fresh samples the specimens are difficult to find, because they are often masked by detritus. In my experience it is best to create microaquaria with high layer thickness and fill them with detritus containing samples. After about 1-2 days

specimens can be found outside the detritus. Either isolate these specimens or try to reduce the layer thickness of the microaquariums as far as the foreign bodies under the coverslip allow.

The division of *Diaphoropodon mobile* occurs along the longitudinal axis (s. fig. 4). The division begins at the posterior end. Not only does the protoplast divide, but the flexible test constricts and also divides.

More images and information on *Diaphoropodon mobile*: [Ferry Siemensma-Microworld-Diaphoropodon mobile](#)




Fig. 1: *Diaphoropodon mobile*. L = 124 μ m. A specimen with extended filopodia and 16–18 μ m long rods (RO). Nu = nucleus. Obj. 40 X.



Fig. 2: *Diaphoropodon mobile*. L = 138 μ m. The test of this specimen is mainly covered with organic material. The rods (RO) are 9–10 μ m long. Obj. 40 X.




Fig. 3 a-b: *Diaphoropodon mobile*. L = 145 μm . Two focal planes of a specimen with extended and branched filopodia (Fi). The rods (arrow) of this specimen are only 4–5 μm long. Obj. 60 X.




Fig. 4: *Diaphoropodon mobile*. L = 65 μm . A specimen during cell division along the longitudinal axis. Obj. 40 X.