

Dicranophoroides caudatus

Ehrenberg, 1834

Most likely ID: n.a.


Synonym: *Dicranophorus caudatus*

Sampling location: [Simmelried](#), [Ulmisried](#)

Phylogenetic tree: [*Dicranophoroides caudatus*](#)

Diagnosis:

- length 180–310 µm
- two frontal eyespots with lenses.
- long, straight toes with pointed ends
- colored green by symbiotic algae (kleptoplasts)
- lacks a retrocerebral sac
- trophi large, rami tips with small teeth



1, 2 = after Wulfert
 3, 4 = after Hudson & Gosse

Dicranophoroides caudatus

Dicranophoroides caudatus is very common in my localities [Simmelried](#) and [Ulmisried](#) and is always found in detritus. There it feeds on Euglenophyceae, like *Euglena*, *Lepocinclus* or *Phacus* (s. fig. 4). The species is very easy to identify by its green coloration. This is caused by so-called kleptoplasts. These are “stolen” chloroplasts that come from the prey organisms (s. fig. 5). The chloroplasts are not digested like the rest of the prey, but are stored intracellularly while retaining their full function. The selection of chloroplasts appears to be selective. I have only been able to find disc-shaped chloroplasts in *Dicranophoroides caudatus*, such as those found in *Lepocinclus acus*. In starving specimens the number of kleptoplasts can be strongly reduced up to almost colorless specimens.

More images and information on *Dicranophoroides caudatus*: [Michael Plewka-Freshwater life-Dicranophoroides caudatus](#)

Dicranophoroides caudatus
Obj. 40 X




Fig. 1 a-b: *Dicranophoroides caudatus*. L = 240 μ m. A freely swimming, well fed specimen in lateral view from right. The stomach is completely filled with disc-shaped paramylon grains (PG), most likely from *Phacus* species. Obj. 40 X.

Dicranophoroides caudatus
Obj. 100 X





Fig. 2 a-b: *Dicranophoroides caudatus*. L = 264 µm. A slightly squashed specimen in detail. CB = circumapical band of cilia. Obj. 100 X.

Dicranophoroides caudatus
Obj. 100 X



© Martin Kreutz

Fig. 3: *Dicranophoroides caudatus*. L = 264 μm . In this squashed specimen, the rami protrude from the mouth with the teeth at the distal ends. RA = rami with distal teeth. Obj. 100 X.

Dicranophoroides caudatus
Obj. 60 X




Fig. 4: *Dicranophoroides caudatus*. In this squashed specimen the paramylon grains (PG) of ingested Euglenophyceae (likely *Lepocinclis acus*) are visible. Obj. 60

X.




Fig. 5: *Dicranophoroides caudatus*. The green coloration is caused by chloroplasts derived from phagocytosed Euglenophyceae (e.g. *Euglena*, *Lepocinclis* or *Phacus*). The chloroplasts of the prey are not digested but are stored intracellularly and maintained in full function. They are then referred to as kleptoplasts (KLP). ES = eye spot. Obj. 100 X.