

***Dicranophorus luetkeni* (Bergendal, 1892)**

Most likely ID: n.a.


Synonym: n. a.

Sampling location: [Simmelried](#)

Phylogenetic tree: [Dicranophorus luetkeni](#)

Diagnosis:

- body slender, elongate and with straight ventral side
- length 180–317 μm
- corona ventral, a little shorter than head
- large rostrum
- with retrocerebral organ
- no eyespot
- toes long, straight with slender, blunt claws
- the claws can be bent



Dicranophorus luetkeni

I have found *Dicranophorus luetkeni* only rarely and exclusively in the [Simmelried](#). The few finds are from October 2006, March 2008 and May 2010, after which I have not found any more specimens.

I recognize the species mainly by the long toes, whose claws can be bent (s. fig. 3). This can be observed especially in slightly squashed specimens. In addition, *Dicranophorus luetkeni* has an apical rostrum above the corona (s. fig. 2d). The corona is limited to the ventral side. The stomach is usually yellow or orange in color.

More images and information on *Dicranophorus luetkeni*: [Michael Plewka-Freshwater life-Dicranophorus luetkeni](#)




Fig. 1 a-d: *Dicranophorus luetkeni*. L = 250 μ m. Lateral view from right of a freely swimming specimen. Obj. 40 X.




Fig. 2 a-d: *Dicranophorus luetkeni*. L = 295 μ m. Lateral view from right of a slightly squashed specimen. Note the apical rostrum (RO). Obj. 40 X.



Fig. 3: *Dicranophorus luetkeni*. L = 295 μm . The same specimen as shown in fig. 2 a-d with focal plane on the bent claws (arrow) of the toes. Obj. 60 X.




Fig. 4: *Dicranophorus luetkeni*. L = 295 μ m. Two focal planes of the trophi in a strongly squashed specimen. Obj. 100 X.