

Euastrum neosinuosum

O.V. Anissimova & Guiry, 2021

Most likely ID: n.a.


Synonym: *Euastrum sinuosum*, *Euastrum circulare* var. *falaisense*

Sampling location: [Schwemm Moor \(Austria\)](#)

Phylogenetic tree: [Euastrum neosinuosum](#)

Diagnosis:

- semi-cells almost trapezoidal with rounded borders
- length 70-85 µm, width 35-45 µm
- apical lobe separated by deep incision
- two lateral lobes and two basal lobes per semi-cell
- semi-cells with each 5 protuberances
- one central pore in each semi-cell
- protuberances covered with inconspicuous warts
- deep, linear sinus



Euastrum neosinuosum

So far, I have only found *Euastrum neosinuosum* in the [Schwemm Moor](#) in Austria. There were a large number of specimens in the samples.

Euastrum neosinuosum can be identified by its trapezoidal semi-cells with lateral, hump-shaped lobes. The apical lobe is divided by a deep incision, which is closed at the top. The cell wall is significantly thickened at the projections and covered with fine warts. Each semi-cell has a total of 5 protuberances and a central pore (s. fig. 1 a-c).

The similar species *Euastrum aboense* is slightly smaller (max. 60 µm), has less pronounced lateral lobes, and has no warts on the surface. In addition, *Euastrum aboense* has 6 more pores between the protuberances of the semi-cells in addition to the central pore.




Fig. 1 a-c: *Euastrum neosinuosum*. L = 85 μ m. Three focal planes of a specimen found in the [Schwemm Moor](#). Each semi-cell has a central pore (PO) and 5 protuberances (1-5). Obj. 100 X.




Fig. 2 a-c: *Euastrum neosinuosum*. L = 85 μm . The same specimen as shown in fig. 1 a-c in brightfield illumination. Obj. 100 X.