

## ***Eudorina elegans* Ehrenberg, 1832**

**Most likely ID:** n.a.


**Synonym:** n.a.

**Sampling location:** [Simmelried](#)

**Phylogenetic tree:** [Eudorina elegans](#)

**Diagnosis:**

- colony ellipsoidal or spherical with a mucilaginous envelope
- length 50–200 µm
- colony consisting of 4–8–16–32–64 cells (in most cases 32)
- cells arranged in 5 layers (4–8–8–8–4)
- cells spherical, sub-spherical or pear-shaped each with 2 flagella of equal length
- flagella cross the mucilaginous envelope through canals
- each cell with 2 contractile vacuoles
- one chloroplast, cup-shaped
- one pyrenoid
- one eyespot




after Hartmann


### Eudorina elegans

*Eudorina elegans* is a widespread volvococcal alga, which sometimes occurs in masses especially in my sampling site [Simmelried](#). The spherical colonies with mostly 32 cells are easy to identify (s. figs. 1 a-c and 2 a-b).

*Eudorina elegans* can reproduce asexually by vegetative division, but also by sexual reproduction, as there are male as well as female colonies. In the vegetative state the sexes cannot be distinguished. Only at stages of reproduction it is possible to determine which sex is present. In the male colonies clusters of sperm cells form (s. figs. 5 and 6) and in the female colonies either immobile aplanospores or flagellated zoospores are formed after fertilization. When immobile aplanospores are formed, they begin to germinate and form new colonies by cell division (s. fig. 7).



**Fig. 1 a-c:** *Eudorina elegans*. D = 132  $\mu\text{m}$  (of colony). The focal planes of a freely swimming colony. Note the eyespots (ES) of the cells. F = flagella, ME = mucilaginous envelope. Obj. 60 X.




**Fig. 2 a-b:** *Eudorina elegans*. D = 98  $\mu\text{m}$  (of colony). A second freely swimming colony. F = flagella, ES = eyespots. Obj. 60 X.




**Fig. 3:** *Eudorina elegans*. D = 131 µm (of colony). The cells of a colony in detail. Note the contractile vacuoles (CV) and the eyespot (ES) of some specialized cells. Obj. 100 X.

*Eudorina elegans*  
Obj. 100 X




**Fig. 4:** *Eudorina elegans*. D = 131 µm (of colony). Focal plane on cells with a visible pyrenoid (PY) and the centrally located nucleus (Nu). Obj. 100 X.



© Martin Kreutz

**Fig. 5:** *Eudorina elegans*. The formation of clusters of sperm cells (CSC) in a male colony. MME = mucilaginous envelope of mother cell. Obj. 40 X.


*Eudorina elegans*  
Obj. 100 X



© Martin Kreutz

**Fig. 6:** *Eudorina elegans*. The released clusters of sperm cells. Obj. 100 X.

*Eudorina elegans*  
Obj. 40 X



**Fig. 7:** *Eudorina elegans*. Aplanospores (AS) and germinating aplanospores (GAS) in a female colony. Obj. 40 X.