## Glaucoma frontata (Stokes, 1886)

Most likely ID: n.a.

Synonym: Dallasia frontata, Diplomastax frontata

Sampling location: Simmelried

Phylogenetic tree: **Glaucoma frontata** 

## **Diagnosis**:

- length about 100–160  $\mu m$
- body is ovoid, spindle-shaped or "tailed" at the posterior end
- oral aperture located in the anterior third is from "Glaucoma"-type with three distinct membranelles
- CV central
- macronucleus globular or oval, centrally located or in the anterior third
- inconspicuous fringe of extrusomes present



after Kahl Glaucoma frontata

I found *Glaucoma frontata* for the first time in <u>Simmelried</u> in 1999. After that this species occured in intervals of several years. The identification of *Glaucoma frontata* according to the drawing and description of Kahl is difficult, because the shape of *Glaucoma frontata* can vary very much. There are varieties from slender spindle-shaped to broadly oval or with a distinctly tapered posterior end. Therefore, I provide some examples of freely swimming specimens with different shapes found between 2008 and 2019 (s. figs. 1 a-b to 6 a-d).



Fig. 1 a-b: *Glaucoma frontata*.  $L = 140 \mu m$ . Freely swimming specimen with a slightly tapered posterior end. Obj. 40 X.



**Fig. 2 a-b:** *Glaucoma frontata*.  $L = 118 \mu m$ . Freely swimming specimen with an almost ovoid shape. Obj. 40 X.



**Fig. 3 a-d:** Glaucoma frontata.  $L = 133 \mu m$ . A freely swimming, spindle-shaped specimen. Obj. 60 X.



Fig. 4 a-d: Glaucoma frontata. L = 120  $\mu$ m. Freely swimming specimen with a distinct "tail". Obj. 40 X.



Fig. 5 a-d: *Glaucoma frontata*.  $L = 160 \mu m$ . A second freely swimming specimen with a distinct "tail". Obj. 100 X.



**Fig. 6 a-d:** Glaucoma frontata.  $L = 160 \mu m$ . A third freely swimming specimen with a distinct "tail". Obj. 100 X.

In a slightly pressed specimen, the structure of the oral aperture can be seen in more detail. The structure is of the typical *Glaucoma* style. Three adoral membranelles (M1 – M3) are parallel and surrounded by an undulating membrane from the right side (s. fig. 8 a-b). The middle adoral membranelle (M2) is most pronounced, just as Corliss drew it (s. fig. 9).





**Fig. 7 a-b:** *Glaucoma frontata*.  $L = 145 \mu m$ . Ventral view (a) and lateral view (b) of a slightly squashed specimen. Ma = macronucleus. Obj. 100 X.





**Fig. 8 a-b:** *Glaucoma frontata*.  $L = 145 \mu m$ . Oral aperture in two focal planes. M1 - M3 = adoral membranelles, UM = undulating membrane. Obj. 100 X.

The ciliation of the oral aperture of *Glaucoma frontata* was described in detail by Corliss (s. fig. 9 a). According to his investigation, a small patch of cilia is located at the anterior end of the prominent adorale membranelle M2, which he called "enigmatic structure" or "X-body".

He suggested that it was a separate piece of the membranelle M2. This structure is difficult to see without silver impregnation. However, when the oral aperture is focussed from the dorsal side of the ciliate, it can be observed (s. fig. 9 b):



**Fig. 9 a-b:** *Glaucoma frontata*. a) the arrangement of the adoral membranelles (M1 – M3) and undulating membrane (UM) of the oral aperture according to Corliss including the enigmatic X-Body. b) focus on the oral aperture of *Glaucoma frontata* from dorsal. M1 – M 3 = adoral membranelles, UM = undulating membrane, X = X-body. Obj. 100 X.