

***Kerona pediculus***  
**(Müller, 1773) Müller, 1786**

**Most likely ID:** n.a.


**Synonym:** *Kerona polyporum*

**Sampling location:** [Pond of the waste disposal company Constance](#)

**Phylogenetic tree:** [Kerona pediculus](#)

**Diagnosis:**

- body kidney-shaped, dorso-ventrally flattened
- length 130–205 µm
- adoral zone about 50 % of body length
- two ellipsoidal macronuclei
- each macronucleus with a spherical micronucleus, clearly separated
- contractile vacuole on left margin, below mouth
- 6 slightly curved frontoventral rows of cirri
- 4–6 buccal cirri
- 5 transverse cirri
- 3 caudal cirri




## Kerona pediculus

*Kerona pediculus* is an ectocommensal ciliate, living on various *Hydra* species, and creeps on them rapidly. The species is feeding on algae and ectodermal cells from *Hydra*.


Because it lives on *Hydra*, *Kerona pediculus* cannot be confused with any other species. The body is kidney-shaped and has a complicated ciliature. However, the diagonal rows of frontoventral cirri and the transvers cirri are easily recognizable (s. fig. 1 b). The three caudal cirri are difficult to recognize and I was also unable to clearly identify the buccal cirri.

*Kerona pediculus* has two ellipsoidal macronuclei. Each macronucleus is associated with a spherical micronucleus, which is clearly separated from the macronucleus (s. fig. 2 b). The adoral zone extends approximately to the middle of the body (s. fig. 2 a).

More images and information on *Kerona pediculus*: [closterium\\_mysterium-iNaturalist-Kerona pediculus](#)



**Fig. 1 a-b:** *Kerona pediculus*. L = 134  $\mu$ m. Two focal planes from ventral. The diagonal running frontoventral cirri (FVC) are visible as well as two of the 5 transverse cirri (TC). Obj. 60 X.



**Fig. 2 a-b:** *Kerona pediculus*. L = 134  $\mu$ m. Focal plane on the adoral zone of membranelles (AZM) and the two macronuclei (Ma 1, Ma 2) with two, clearly separated micronuclei (Mi 1, Mi 2). Obj. 60 X.