

***Lepadella rhomboides* (Gosse, 1886)**

Most likely ID: n.a.


Synonym: *Lepadella rhomboides* var. *carinata*, *Lepadella rhomboides* var. *haueri*

Sampling location: [Simmelried](#)

Phylogenetic tree: [Lepadella rhomboides](#)

Diagnosis:

- lorica ovoid to rhomboid-ovate
- length of lorica 110–120 µm, width 55–88 µm
- dorsally with a moderately high keel, sides of keel convex
- ventral side flat with delicate longitudinal folds
- ventral sinus V-shaped
- foot groove U-shaped
- three foot segments, terminal foot segment longer than basal section of foot
- slender toes pointed
- two eyespots



after Harring

Lepadella rhomboides

I have found *Lepadella rhomboides* so far exclusively in the [Simmelried](#). However, the species is not very common. I documented findings in June 2004 and in July 2008. I recognize *Lepadella rhomboides* by the slender, oval shape, the V-shaped ventral sinus (s. fig. 1a) and the fine longitudinal folds on the ventral side of the lorica (s. fig. 3).

Lepadella rhomboides
Obj. 40 X




Fig. 1 a-b: *Lepadella rhomboides*. L = 120 μ m (of the lorica). Ventral view of a slightly squashed specimen. Note the V-shaped sinus (VS) and the elongated terminal foot segment (TF). Obj. 40 X.



Fig. 2 a-b: *Lepadella rhomboides*. L = 115 μm (of the lorica). Two focal planes from dorsal. Note the ridge of the dorsal keel (DK). Vit = vitellarium. Obj. 40 X.



Fig. 3: *Lepadella rhomboides*. L = 85 μm (of the lorica). Ventral view of a slightly squashed specimen. Note the delicate longitudinal folds of the lorical (LF). Obj. 100 X.




Fig. 4 a-c: *Lepadella rhomboides*. L = 127 μm (of the lorica). Ventral view of a slightly squashed specimen found in November 2022. Obj. 60 X.



Fig. 5: *Lepadella rhomboides*. L = 115 μm (of the lorica). Lateral view from left of a freely swimming specimen. Obj. 40 X.