

## ***Lepadella rhomboides* Gosse, 1886**

**Most likely ID:** n.a.


**Synonym:** *Lepadella rhomboides* var. *carinata*, *Lepadella rhomboides* var. *haueri*

**Sampling location:** [Simmelried](#)

**Phylogenetic tree:** [Lepadella rhomboides](#)

### **Diagnosis:**

- lorica ovoid to rhomboid-ovate
- length of lorica 110–120 µm, width 55–88 µm
- dorsally with a moderately high keel, sides of keel convex
- ventral side flat with delicate longitudinal folds
- ventral sinus V-shaped
- foot groove U-shaped
- three foot segments, terminal foot segment longer than basal section of foot
- slender toes pointed
- two eyespots



after Harring

### Lepadella rhomboides

I have found *Lepadella rhomboides* so far exclusively in the [Simmelried](#). However, the species is not very common. I documented findings in June 2004 and in July 2008. I recognize *Lepadella rhomboides* by the slender, oval shape, the V-shaped ventral sinus (s. fig. 1a) and the fine longitudinal folds on the ventral side of the lorica (s. fig. 3).

*Lepadella rhomboides*  
Obj. 40 X



**Fig. 1 a-b:** *Lepadella rhomboides*. L = 120  $\mu$ m (of the lorica). Ventral view of a slightly squashed specimen. Note the V-shaped sinus (VS) and the elongated terminal foot segment (TF). Obj. 40 X.

*Lepadella rhomboides*  
Obj. 40 X




**Fig. 2 a-b:** *Lepadella rhomboides*. L = 115  $\mu$ m (of the lorica). Two focal planes from dorsal. Note the ridge of the dorsal keel (DK). Vit = vitellarium. Obj. 40 X.



**Fig. 3:** *Lepadella rhomboides*. L = 85  $\mu$ m (of the lorica). Ventral view of a slightly squashed specimen. Note the delicate longitudinal folds of the lorical (LF). Obj. 100



X.



**Fig. 4 a-c:** *Lepadella rhomboides*. L = 127  $\mu\text{m}$  (of the lorica). Ventral view of a slightly squashed specimen found in November 2022. Obj. 60 X.



**Fig. 5:** *Lepadella rhomboides*. L = 115  $\mu\text{m}$  (of the lorica). Lateral view from left of a freely swimming specimen. Obj. 40 X.