Litonotus cygnus Müller, 1773

Most likely ID: n.a.


Synonym: n.a.

Sampling location: Simmelried, Purren pond, Mainau pond, Bussenried, Buendtlisried, Ulmisried

Phylogenetic tree: Litonotus cyanus

Diagnosis:

- body slenderly lanceolate and flattened, ventral side convex, dorsal side straight
- posterior end norrowly rounded or tail-like
- neck stretched more than body length, bent dorsally
- length 200-300 µm
- left side with some rows of short bristles
- right side with 6-13 rows of cilia
- dorsal brush of club-shaped, short cilia
- two spherical macronuclei
- one spherical micronucleus between the macronuclei
- extrusomes rod-shaped, slightly curved, 6-8 µm long
- extrusomes arranged mainly along oral cleft
- contractile subterminal, at the end of a left-sided bulge
- sometimes a second contractile vacuole in anterior third



Litonotus cygnus

I find *Litonotus cygnus* in almost all of my sampling sites. The ciliate is difficult to observe in fresh specimens, but it likes to settle on the <u>floating coverslip</u>. Litonotus cygnus then glides along it with its right, ciliated side.

Litonotus cygnus is easy to identify. It is the largest of all *Litonotus* species. Typical are the two spherical macronuclei, which can be recognized even at low magnifications. The neck is very long and, especially at the front end, is clearly bent backwards and slightly widened. The posterior end is almost always tail-shaped in my population. I have only rarely found narrowly rounded forms. The extrusomes are found almost exclusively on the ventral side of the neck. This is where the oral cleft is located, but it is only visible during feeding.

In rare cases, in addition to the subterminal contractile vacuole, a second, smaller one is visible in the anterior third (s. fig. 3). It is less active than the subterminal contractile vacuole and is particularly visible in slightly compressed specimens.




Fig. 1 a-d: Litonotus cygnus. L = 180 μm . Different focal planes of a freely swimming

specimen from right. CV = contractile vacuole, Ma = two spherical parts of the macronucleus. Obj. 60 X.




Fig. 2 a-d: Litonotus cygnus. $L=180~\mu m$. A second specimen from right (a-c) and with focal planes on the left side (d). Obj. 60 X.




Fig. 3: Litonotus cygnus. $L=180~\mu m$. The slightly specimen as shown in fig. 2 a-d. Note the second contractile vacuole (CV 1) in the anterior third. CV 2 = subterminal contractile vacuole. Obj. 60 X.




Fig. 4 a-b: Litonotus cygnus. The anterior end with the fringe of extrusomes (EX) and the dosal brush (DB) in detail. The extrusomes are curved and 6.5 μ m long. Obj. 100 X.

