

***Metopus nasutus* Da Cunha, 2015**

Most likely ID: n.a.

Synonyms: *Copemetopus nasutus*, *Metopus vexilliger*

Sampling location: [Purren pond](#), [Simmelried](#)

Phylogenetic tree: [Metopus nasutus](#)

Diagnosis:

- body elongated oval, posterior half slightly tapered, dorso-ventrally slightly flattened
- apical end elongated to a trunk-shaped organelle with pointed distal end, 35–50 µm long
- trunk-shaped organelle towards the left side of the body
- length 100–130 µm
- adoral zone reaches the cell equator
- pharynx slightly below cell equator, bent left
- macronucleus spherical or ellipsoidal with one adjacent, spherical micronucleus
- fringe of about 1 µm long extrusomes beneath the pellicle
- contractile vacuole terminal
- caudal cilia 10–30 µm long



after Kahl

Metopus nasutus

Metopus nasutus is one of the most common metopid ciliates in the [Simmelried](#) and the [Purren pond](#). Especially in the [Purren pond](#) this remarkably shaped *Metopus* species can be found reliably. *Metopus nasutus* is easily recognized by a trunk-shaped organelle located at the anterior end. This organelle of unknown function makes the species unmistakable.

In my population I could find specimens with body lengths between 90–170 μm , which extends the range given by Kahl (100–150 μm). The trunk-shaped organelle is passively dragged along when swimming and lies against the left side of the body, as Kahl describes it (s. fig. 7 a-b). Only once could I observe the organelle to be stretched forward (s. fig. 3 a-d). In this specimen the trunk-shaped organelle was also unusually long with 74 μm . Kahl gives a length of 10–30 μm . On which factors the length of the organelle depends is unknown. In any case, it can melt under mechanical stress (coverslip pressure).

At high magnification, many vesicles and granules can be seen in the trunk-shaped organelle (s. fig. 6). In one case, I was able to find an attached bacterium at the tip of the organelle (s. fig. 6). Possibly this is a coincidental constellation, however it made me think of the possibility that the trunk-shaped organelle may be used like a glue rod to collect bacteria and other food particles. This theory is supported by the observation that in some free-swimming specimens I had the impression that the trunk was sometimes located between the perizonal cilia and the adoral membranelles in the direction of the mouth opening (s. figs. 8 a-b and fig. 9). Possibly bacteria or other food particles are „brushed off“ between the perizonal cilia and the adoral membranelles during this behavior.

Beneath the pellicle of *Metopus nasutus* a fringe of extrusomes about 1 µm long is located (s. fig. 5a), not mentioned by Kahl. Many symbiotic bacteria are visible in the cytoplasm (s. fig. 5 a), especially in the area around the macronucleus. In the food vacuoles I found colorless bacteria, green-yellow sulfur bacteria, pink rhodobacteria and small green algae.




Fig. 1 a-c: *Metopus nasutus*. L = 165 μ m. Ventral view of a freely swimming specimen. Note the trunk-shaped organelle (TSO) at apical end. AZM = adoral zone of membranelles, CC = caudal cilia, CV = contractile vacuole, MO = mouth opening. Obj. 60 X.



Fig. 2 a-b: *Metopus nasutus*. L = 115 μ m. Ventral view of a seond, freely swimming specimen. CC = caudal cilia, PC = perizonal cilia, TSO = trunk-shaped organelle. Obj. 60 X.




Fig. 3 a-d: *Metopus nasutus*. L = 122 µm (without trunk shaped organelle). A freely swimming specimen with an exceptionally long trunk-shaped organelle (L = 74 µm) that is extended forward in the direction of swimming. CC = caudal cilia, CV = contractile vacuole, MO = mouth opening. Obj. 40 X.




Fig. 4 a-c: *Metopus nasutus*. L = 92 μ m. Focal plane on the trunk-shaped organelle (TSO) in an freely swimming specimen. Note the long perizonal cilia (PC). Obj. 60 X.




Fig. 5 a-b: *Metopus nasutus*. A slightly squashed specimen from ventral (a) and dorsal (b). Note the symbiotic bacteria in the cytoplasm (SB) and the distinct striation of the granulated pellicle (b). Beneath the pellicle a fringe of about 1 μm

long extrusomes is visible (FE): Ma = macronucleus, Mi = micronucleus, MO = mouth opening, TSO = trunk-shaped organelle. Obj. 100 X.

Metopus nasutus
Obj. 100 X

BAC

10 μ m

TSO

© Martin Kreutz

Fig. 6: *Metopus nasutus*. The trunk-shaped organelle (TSO) in detail. In the trunk many vesicles and granules are located. At the tip of the trunk an adherent bacterium is visible (BAC). This may be there by chance or it may be an indication that the trunk has a function similar to a glue rod to collect bacteria. Obj. 100 X.



Fig. 7 a-b: *Metopus nasutus*. L = 115 μ m. A freely swimming specimen from left (a) and from dorsal (b). The trunk-shaped organelle (TSO) lies passively on the left side

of the body when swimming. AZM = adoral zone of membranelles, CC = caudal cilia, CV = contractile vacuole, L/R = left/right, Ma = macronucleus, Mi = micronucleus, PCR = perizonal cilia row. Obj. 60 X.




Fig. 8 a-b: *Metopus nasutus*. L = 96 μ m. A freely swimming specimen from left with the trunk-shaped organelle (TSO) bent in the direction of the mouth opening. Obj. 60 X.



Fig. 9: *Metopus nasutus*. L = 148 μ m. In this freely swimming specimen the trunk-shaped organelle (TSO) seems to be located between the perizonal cilia row and the adoral zone, bent in direction of the mouth opening (MO). Obj. 40 X.