

Monactinus simplex var. echinulatum

(Wittrock) Pérez, Maidana & Comas, 2009

Most likely ID: n.a.


Synonym: *Pediastrum simplex var. echinulatum*, *Pediastrum sturmii var. echinulatum*

Sampling location: [Pond of the waste disposal company Constance](#)

Phylogenetic tree: [Monactinus simplex var. echinulatum](#)

Diagnosis:

- coenobium star-shaped, flat and single-layered
- diameter coenobium up to 250 µm
- coenobium of 4, 8, 16, 32, 64, (128) cells
- coenobium covered with distinct warts or short spines
- marginal cells with a single, gradually tapering projection
- inner cells polygonal shaped, concentrically arranged, sometimes with small intercellular spaces
- marginal cells occasionally bearing tufts of mucilaginous spines
- one chloroplast, filling the cell
- single pyrenoid



after Hortobágyi

Monactinus simplex var. *echinulatum*

Monactinus simplex var. *echinulatum* is described as not very common (Komárek & Fott, 1983). However, in the [pond of the waste disposal plant of Constance](#) I found this form in large quantities in the plankton in September 2023.

The genus *Monactinus* is easily recognized because the marginal cells have only one horn-shaped process. In the form *Monactinus simplex* var. *echinulatum*, the gaps between the inner cells are either absent or very small. As a distinctive characteristic, the cell walls of the coenobia are covered with distinct warts or short spines. In my population these were short spines of 0.7 – 1.2 µm length (s. fig. 1 a-b).




Fig. 1 a-b: *Monactinus simplex* var. *echinulatum*. $D = 65\ \mu\text{m}$. Two focal planes of a coenobium of 16 cells. The coenobium is covered with distinct, short spines (arrows) with a length of $0.7 - 1.2\ \mu\text{m}$. Note the delicate mucilaginous spines (MS) arising from the marginal cells. PY = pyrenoid. Obj. 100 X.