

***Nephrocytium limneticum* Smith, 1933**

Most likely ID: n.a.


Synonym: *Gloeocystopsis limnetica*

Sampling location: [Hagstaffel pond](#)

Phylogenetic tree: [Nephrocytium limneticum](#)

Diagnosis:


- coenobia of 4-8-16 cells in mucilage envelope
- daughter colonies 20-50 µm in diameter
- daughter cells remain united within gelatinized cell wall of mother cell
- cells 10-20 µm long, kidney-shaped, slightly curved with rounded apices
- one pyrenoid
- one parietal chloroplast
- planctonic lifestyle



after Guarrera

Nephrocytium limneticum

I find *Nephrocytium limneticum* very rarely in plankton samples, always with several years between them. *Nephrocytium* can be confused with *Kirchneriella*, whose cells are also kidney-shaped or crescent-shaped. However, the colonies of *Nephrocytium* are covered by a distinct and thick layer of mucus, which is absent in *Kirchneriella*. In addition, the daughter colonies of *Nephrocytium* remain united for a long time in the swollen and gelatinized cell wall of the mother cell (s. fig. 2 a).



50 μm

© Martin Kreutz

Fig. 1: *Nephrocytium limneticum*. L = 156 (of coenobium). A coenobium of 8 colonies with 16 cells each within the gelatinized cell wall of the mother cell. Obj. 40 X.

Nephrocytium limneticum
Obj. 100 X

ME

RMC

a

PY

Nu

10 μ m

PY

b

© Martin Kreutz

Fig. 2 a-b: *Nephrocytium limneticum*. L = 10-12 μm (of the cells). A slightly squashed (a) and a strongly squashed (b) coenobium of 4 colonies with 4 cells each within the remains of the mother cell (RMC). Each colony is in a thick mucilage envelope (ME). Nu = nucleus, PY = pyrenoid. Obj. 100 X.




Fig. 3 a-b: *Nephrocytium limneticum*. L = 19-21 μm (of the cells). Two focal planes of a second coenobium of 8 cells. Obj. 100 X.