

***Peloploca fibrata* (Skuja, 1956)**

Most likely ID: n.a.


Synonym: n.a.

Sampling location: [Simmelried](#)

Phylogenetic tree: [Peloploca fibrata](#)

Diagnosis:

- aggregates of cells are multilayered bundles of filaments up to 1 mm long
- ends of the aggregates irregularly tapered or defibered
- single bacteria cells are arranged tightly in the long filaments
- each filament with an indistinct hyaline sheath
- segmentation of the cylindrical single cells in the filaments indistinct, sometimes not visible
- length of cells variable, up to 5 times longer than wide
- cells hyaline, without gas vacuoles



after Skuja

Peloploca fibrata

Peloploca fibrata is an aggregate-forming bacterium that was first described in 1956 by Skuja in Swedish Lapland. In the aggregates, which can consist of more than 1000

individual cells, the bacteria are arranged in filaments in chains. These filaments in turn are arranged in bundles of varying thickness and length to form the aggregate (s. figs. 1 a-b, 2 and 3).

I find *Peloploca fibrata* regularly, but never frequently in the [Simmelried](#). I have not yet been able to find this bacterium in my other sampling sites. *Peloploca fibrata* is almost always associated with rhodobacteria. Since the aggregates of *Peloploca fibrata* are very long (up to 1 mm), they are immediately noticeable in the samples even at low magnifications. In the aggregates of my population, the individual cells in the filaments were not or only indistinctly recognizable, as Skuja has already described. I only found a few aggregates in which the segmentation into individual cells could be clearly seen (s. figs. 4 and 5). They were almost always aggregates with only a few filaments. The individual cells may be easier to recognize in the growth phase of the aggregates. In such aggregates it can be seen that the cylindrical bacteria in the filaments have very different lengths, which is also described by Skuja. Important characteristics of *Peloploca fibrata* are that there are no constrictions between the individual cells and that the cells do not contain gas vacuoles. This distinguishes *Peloploca fibrata* from [Peloploca taeniata](#).

Peloploca fibrata
Obj. 100 X

25 µm

© Martin Kreutz

a

b

Fig. 1 a-b: *Peloploca fibrata*. L = about 500 μm (of the aggregate). Two focal planes of a slightly twisted aggregate of about 20-30 filaments. The segmentation of the single bacteria cells is only visible in few parts of the aggregate. Obj. 100 X.




Fig. 2: *Peloploca fibrata*. L = about 600 μm (of the aggregate). A section of a second aggregate. Obj. 100 X.




Fig. 3: *Peloploca fibrata*. L = about 500 μm (of the aggregate). A section of a third aggregate. Obj. 100 X.



Fig. 4: *Peloploca fibrata*. L = about 300 μm (of the aggregate). In this aggregate of about 10 filaments the single bacteria cells in the filaments were clearly separated. Probably the separation of the cells is better visible at the beginning of the aggregate formation. Obj. 100 X.



Fig. 5: *Peloploca fibrata*. In this section of the fig. 4 the different lengths of the bacteria in the filaments becomes visible. Note that the cells are hyaline without gas vacuoles. Obj. 100 X.