

***Peloploca taeniata* Lauterborn, 1913**

Most likely ID: n.a.


Synonym: n.a.

Sampling location: [Simmelried](#)

Phylogenetic tree: [Peloploca taeniata](#)

Diagnosis:

- aggregates of cells are multilayered bundles of filaments up to 1 mm long
- aggregates end in single filaments
- single bacteria cells are arranged tightly in the long filaments
- filaments without hyaline sheath, aggregates with 5 µm thick hyaline sheath
- cross walls between cells slightly constricted, hard to see
- single cells 0.7-1 µm x 3-10 µm with distinct gas vacuoles



after Skuja

Peloploca taeniata

Peloploca taeniata is an aggregate-forming bacterium that was first described by Lauterborn in 1913. However, his description is very brief and not very informative. His drawings were also rather superficial. It was not until 1956 that Skuja published a re-description of this species in his results regarding the investigation of Swedish inland water bodies. His description is much more precise and his drawings more informative.

The aggregates of *Peloploca taeniata* consist of filaments of bacteria. These filaments in turn are arranged in bundles of varying thickness and length. The aggregates can therefore contain well over 1000 cells.

I find *Peloploca taeniata* regularly, but never frequently in the [Simmelried](#). I have not yet been able to find this bacterium in my other localities. *Peloploca taeniata* is almost always associated with rhodobacteria (s. fig. 1). Since the aggregates of *Peloploca taeniata* are very long (up to 1 mm), they are immediately noticeable in the specimens even at low magnifications. The individual cells in the filaments were only clearly distinguishable in strongly squashed aggregates (s. fig. 4). According to my measurements, the individual cells are between 5–7 μm long, which fits well with the measurements of Skuja, who states 3–10 μm . Each cell contains one or two gas vacuoles, which gives the aggregates an almost glittering appearance. I often find the similar species [Peloploca fibrata](#) co-occurring with *Peloploca taeniata* in the samples. However, it is easy to distinguish the two species because [Peloploca fibrata](#) has no gas vacuoles and the cells are clear and hyaline.



Fig. 1: *Peloploca taeniata*. L = about 600 μm (of aggregate). An aggregate, consisting of a

bundle of filaments, between other bacteria and rhodobacteria. Obj. 100 X.

Peloploca taeniata
Obj. 100 X

25 μm

© Martin Kreutz

Fig. 2: *Peloploca taeniata*. L = about 700 μm (of aggregate). A slightly twisted aggregate. Due to the gas vacuoles in the cells the aggregate appears „glittering“. Obj. 100 X.




Fig. 3: *Peloploca taeniata*. L = about 500 μm (of aggregate). A more wavy and ribbon-shaped aggregate. Obj. 100 X.




Fig. 4: *Peloploca taeniata*. A severed piece of a squashed aggregate. Note the gas vacuoles (GV) in the cells and the slight constrictions of the cross walls (CCW) between the cells. Obj.

100 X.