

***Proalinopsis caudatus* Collins, 1872**

Most likely ID: n.a.


Synonym: n.a.

Sampling location: [Simmelried](#)

Phylogenetic tree: [*Proalinopsis caudatus*](#)

Diagnosis:

- body elongated spindle-shaped, head clearly set off
- dorsal side curved, ventral flat
- length 125–268 µm
- corona reduced, ventrally shifted
- cuticle clear, transparent, covered with mucilaginous sheath
- mucilaginous sheath often interspersed with bacteria
- stomach and intestine clearly separated
- lateral antennae with long setae
- dorsal antenna and caudal antenna on a distinct papilla
- one eyespot with lens
- toes 16–22 µm long, narrow and pointed
- two distinct pedal glands



1 = after Koste
 2, 3 = after Wulfert
 4 = after Harring & Myers

Proalinopsis caudatus

I find *Proalinopsis caudata* regularly in the Simmelried, especially in old samples with decaying plants. This rotifer has a mucilaginous sheath (s. figs. 3 b, 4 a and 5), which can be seen well especially in low magnifications. The head is clearly separated from the body (s. fig. 1 a-b), which can be seen especially well in lateral view. The locomotion is a slow gliding.

The mucilaginous sheath is interspersed with bacteria. However, these are not interspersed randomly, but are radially oriented and are mostly located in the outer layer of the mucilaginous sheath. In my population, the bacteria in the sheath were 2–4 µm long rods (s. fig. 5). Whether the bacteria use the excreted mucilaginous sheath as a food source, or whether they also provide a benefit to *Proalinopsis caudatus*, cannot be readily determined. However, it is striking that they are obviously bacteria of the same species and are aligned along the growth structure of the mucilaginous sheath.

The large rotifer *Notommata copeus* has also got a mucilaginous sheath that is interspersed with bacteria in the same manner. This rotifer also moves only by slowly gliding and is often found between floating plant masses. Obviously the presence of a mucilaginous sheath offers an advantage for this lifestyle, which remains unresolved for the time being.

More images and information on *Proalinopsis caudatus*: [Michael Plewka-Freshwater life-Proalinopsis caudatus](#)




Fig. 1 a-b: *Proalinopsis caudatus*. L = 180 µm. A specimen gliding along a detritus flake from left lateral view. Note the dorsal antenna (DA) and the caudal antenna (CA) arising from short papillae. The head is clearly set off from the body. Obj. 40 X.




Fig. 2 a-c: *Proalinopsis caudatus*. L = 217 μ m. A freely gliding specimen from dorsal. Note the lateral antennae with the long setae. The body is covered with a mucilaginous sheath (MS), visible by interspersed bacteria. Obj. 40 X.

Proalinopsis caudatus
Obj. 40 X



Fig. 3 a-b: *Proalinopsis caudatus*. L = 217 μ m. The slightly squashed specimen shown in fig. 2 a-c. ES = eyespot, GG = gastric glands, Int = intestine, MS = mucilaginous sheath, ST = stomach, TO = toes, Vit = vitellarium. Obj. 40 X.

Proalinopsis caudatus
Obj. 60 X




Fig. 4 a-b: *Proalinopsis caudatus*. L = 216 μ m. A transparent, slightly squashed specimen from ventral. Note the mucilaginous sheath (MS) with the interspersed bacteria. Obj. 60 X.




Fig. 5: *Proalinopsis caudatus*. L = 216 μ m. Focal plane on the surface of the mucilaginous sheath with the interspersed bacteria (arrows). The bacteria are 2-4 μ m long rods and arranged radially. Obj. 100 X.

Proalinopsis caudatus
Obj. 60 X



Fig. 6 a-b: *Proalinopsis caudatus*. L = 140 µm. A freely gliding specimen from ventral. Co = corona, PG = pedal glands. Obj. 60 X.

Proalinopsis caudatus
Obj. 100 X




Fig. 7: *Proalinopsis caudatus*. A slightly squashed specimen with focal plane on the eyespot (ES) with an attached lens (LE). Obj. 100 X.




Fig. 8 a-b: *Proalinopsis caudatus*. Two focal planes of the trophi in a strongly squashed specimen. Obj. 100 X.