Proalinopsis caudatus (Collins, 1872)

Most likely ID: n.a.

Synonym: n.a.

Sampling location: Simmelried

Phylogenetic tree: Proalinopsis caudatus

Diagnosis:

- · body elongated spindle-shaped, head clearly set off
- dorsal side curved, ventral flat
- length 125-268 µm
- corona reduced, ventrally shifted
- cuticle clear, transparent, covered with mucilaginous sheath
- mucilaginous sheath often interspersed with bacteria
- stomach and intenstine clearly separated
- lateral antennae with long setae
- dorsal antenna and caudal antenna on a distinct papilla
- one eyespot with lens
- toes 16–22 μm long, narrow and pointed
- two distinct pedal glands



Proalinopsis caudatus

I find *Proalinopsis caudata* regularly in the Simmelried, especially in old samples with decaying plants. This rotifer has a mucilaginous sheath (s. figs. 3 b, 4 a and 5), which can be seen well especially in small magnifications. The head is clearly separated from the body (s. fig. 1 a-b), which can be seen especially well in lateral view. The locomotion is slowly gliding.

The mucilaginous sheath is interspersed with bacteria. However, these are not interspersed randomly, but are radially oriented and are mostly located in the outer layer of the mucilaginous sheath. In my population, the bacteria in the sheath were 2-4 µm long rods (s. fig. 5). Whether the bacteria use the excreted mucilaginous sheath as a food source, or whether they also provide a benefit to *Proalinopsis caudatus*, cannot be readily determined. However, it is striking that they are obviously bacteria of the same species and are aligned along the growth structure of the mucilaginous sheath.

The large rotifer *Notommata copeus* also has a mucilaginous sheath that is interspersed with bacteria in the same manner. This rotifer also moves only slowly gliding and is often found between floating plant masses. Obviously the presence of a mucilaginous sheath offers an advantage for this lifestyle, which remains unresolved for the time being.

More images and information on *Proalinopsis caudatus*: <u>Michael Plewka-Freshwater life</u>-<u>*Proalinopsis caudatus*</u>



Fig. 1 a-b: *Proalinopsis caudatus.* $L = 180 \mu m$. A specimen gliding along of a detritus flake from left lateral view. Note the dorsal antenna (DA) and the caudal antenna (CA) arising from short papillae. The head is clearly set off from the body. Obj. 40 X.



Fig. 2 a-c: *Proalinopsis caudatus.* $L = 217 \mu m$. A freely gliding specimen from dorsal. Note the lateral antenane with the long setae. The body is covered with a mucilaginous sheath (MS), visible by interspersed bacteria. Obj. 40 X.



Fig. 3 a-b: *Proalinopsis caudatus.* $L = 217 \mu m$. The slightly squashed specimen shown in fig. 2 a-c. ES = eyespot, GG = gastric glands, Int = intestine, MS = mucilaginous sheath, ST = stomach, TO = toes, Vit = vitellarium. Obj. 40 X.



Fig. 4 a-b: Proalinopsis caudatus. $L = 216 \mu m$. A transparent, slightly squashed specimen from ventral. Note the mucilaginous sheat (MS) with the interspersed bacteria. Obj. 60 X.



Fig. 5: *Proalinopsis caudatus*. L = 216 μ m. Focal plane on the surface of the mucilaginous sheath with the interspersed bacteria (arrows). The bacteria are 2–4 μ m long rods and arranged radially. Obj. 100 X.



Fig. 6 a-b: *Proalinopsis caudatus.* $L = 140 \mu m$. A freely gliding specimen from ventral. Co = corona, PG = pedal glands. Obj. 60 X.



Fig. 7: *Proalinopsis caudatus.* A slightly squashed specimen with focal plane on the eyespot (ES) with an attached lens (LE). Obj. 100 X.



Fig. 8 a-b: *Proalinopsis caudatus.* Two focal planes of the trophi in a strongly squashed specimen. Obj. 100 X.