

Pseudovorticella fasciculata

(Müller, 1773) Foissner & Brozek, 1996

Most likely ID: n.a.


Synonym: *Vorticella fasciculata*, *Pseudovorticella chlorelligera*, *Vorticella margaritata* var. *chlorelligera*

Sampling location: [Simmelried](#)

Phylogenetic tree: [Pseudovorticella fasciculata](#)

Diagnosis:

- length 60–70 µm, width 50–55 µm
- cells inverted bell-shaped
- contracted cells broad ovoid, diameter 44–50 µm
- stalk 300–500 µm long, contracts in spirals
- plasm green due to symbiotic algae
- symbiotic algae spherical, diameter 3–4 µm
- two contractile vacuoles
- macronucleus J-shaped
- pellicle covered with pellicular tubercles.
- pellicular tubercles spherical, widely spaced, different sizes
- solitary or in pseudocolonies




Pseudovorticella fasciculata

The species *Pseudovorticella fasciculata* has undergone many name changes and synonyms since its discovery by Müller in 1773. Müller himself placed the species in the to the genus *Vorticella* (*Vorticella fasciculata*). Later, Kahl stated that *Vorticella fasciculata* is apparently identical to the species *Vorticella margaritata* var. *chlorelligera* described by him. Only in 1975 Foissner and Schiffmann assigned this species on the basis of the silver line system to the genus *Pseudovorticella*. The genus *Pseudovorticella* has a latticed silver line system, whereas species of the genus *Vorticella* possess a transverse stripe pattern. Subsequently, Jankowski (1976) renamed *Vorticella margaritata* var. *chlorelligera* to *Pseudovorticella chlorelligera*. Finally, in 1996 Foissner and Brozek re-introduced the original species name *fasciculata* which led to the current name *Pseudovorticella fasciculata*.

There are only few records and descriptions of *Pseudovorticella fasciculata*. I found this species in 1997 in the Simmelried. One could see a green fringe near the surface in the samples with the naked eye, where the pseudocolonies of *Pseudovorticella fasciculata* settled. They showed a clear positive phototactic reaction, because they always collected on the side of the vessel facing the light.

Pseudovorticella fasciculata
Obj. 10 X



© Martin Kreutz




Fig. 1 -2: *Pseudovorticella fasciculata*. L = 65-70 μm . Freely moving specimens of a pseudocolony. Obj. 10 X (fig. 1), Obj. 20 X (fig. 2).





Fig. 3: *Pseudovorticella fasciculata*. L = 65–70 μm . Freely moving specimens of a second pseudocolony. Obj. 10 X.

The cells in my population had a length of 60–70 μm . Kahl gives 60 μm for the synonymous species *Vorticella margaritata* var. *chlorelligera* and Jankowski gives up to 80 μm for the synonymous species *Pseudovorticella chlorelligera*. Due to the symbiotic algae the individuals show an intense green coloration. The body is broadly bell-shaped. A constriction in the posterior third of the body is conspicuous (s. fig. 4). Below this constriction the protoplasm is clear and free of symbiotic algae. In lateral view, this gives the impression of a crescent-shaped, clear plasma fringe. This is a very characteristic feature of *Pseudovorticella fasciculata* and has also been aptly drawn by Kahl in the synonymous species *Vorticella margaritata* var. *chlorelligera* (s. above, drawing 3, after Kahl).



Fig. 4: *Pseudovorticella fasciculata*. L = 63-67 μm . Freely moving specimens of a pseudocolony. Note the constriction at the posterior third of the cell (arrow) and the clear zone of plasm free from symbiotic algae below this constriction. Obj. 40 X.

Pseudovorticella fasciculata
Obj. 40 X



© Martin Kreutz

Fig. 5: *Pseudovorticella fasciculata*. Several contracted cells of a pseudocolony. The focal plane is on the pellicular tubercles. The blister-shaped tubercles are hard to see due to the green background. Obj. 40 X.

The number of contractile vacuoles has not been reported by previous authors. In the squeezed specimens I could clearly see two contractile vacuoles (s. fig. 6). The macronucleus is J-shaped. I could not identify a micronucleus. The symbiotic algae are spherical and very small with a diameter of 3–4 μm . They have a single chloroplast with one pyrenoid and a nucleus (s. fig. 7). The color of the symbiotic algae is green yellowish.




Fig. 6: *Pseudovorticella fasciculata*. In a squashed cell the two contractile vacuoles are visible (CV 1, CV 2). The J-shaped macronucleus (Ma) is labelled by arrows. The symbiotic algae (SA) are very small with a diameter of 3-4 μm . Obj. 100 X.




Fig. 7: *Pseudovorticella fasciculata*. The symbiotic algae (SA) in detail. All symbiotic alga cells have a single chloroplast with a pyrenoid (PY) and a nucleus (Nu). The color of them is green yellowish. Obj 100 X.