## Pyramimonas tetrarhynchus

## Schmarda, 1849

Most likely ID: n.a.


**Synonym:** Pyramidomonas tetrarhynchos

**Sampling location:** Simmelried

Phylogenetic tree: Pyramimonas tetrarhynchos

## **Diagnosis:**

- cells obovid, heart-shaped or inversely pyramidal
- in apical view quadrangular with 4 lobes
- length 18-27 μm
- 4 apical flagella of equal length
- chloroplast cup-shaped, divided apically in 8 lobes
- one pyrenoid, located posteriorly
- one eyespot (on level of pyrenoid)
- one apical contractile vacuole



Pyramimonas tetrarhynchus

So far I have only found *Pyramimonas tetrarhynchus* in the <u>Simmelried</u>, where the species rarely occurs. I usually find specimens in the spring between March and April.

The cells of *Pyramimonas tetrarhynchus* have 4 flagella and 4 longitudinal lobes, which are separated by furrows. This can be seen in apical view. Then the cells appear cloverleafshaped or square with rounded corners (s. drawing 2 above). The eyespot is clearly visible and lies in the posterior third, approximately at the same height as the pyrenoid. The cupshaped chloroplast splits into 4 lobes, which are localized in the 4 longitudinal lobes. Apically, these 4 lobes split again to a total of 8 lobes, which is hard to recognize in living specimens.

Many authors refer to Pyramimonas rhynchomonas as Pyramidomonas rhynchomonas, for example Huber-Pestalozzi (1961).



Fig. 1: Pyramimonas tetrarhynchos.  $L = 21-26 \mu m$ . An aggregation of several, freely swimming specimens. Obj. 100 X.




Fig. 2 a-d: Pyramimonas tetrarhynchos.  $L = 25 \mu m$ . Different focal planes of a freely swimming specimen. Note the 4 apical flagella and the posteriorly located eyespot (ES). Obj. 100 X.




Fig. 3 a-d: Pyramimonas tetrarhynchos.  $L = 26 \mu m$ . A second freely swimming specimen. Note the pyrenoid (PY) on the level of the eyespot (ES). CV = contractile vacuole, Nu = nucleus. Obj. 100 X.