

Rhodobacteria 3

Most likely ID: n.a.

Synonym: n.a

Sampling location: [Simmelried](#)

Phylogenetic tree: n.a.

Diagnosis:

- the cells are oval, larger cells oblong, sometimes irregularly
- length 4 – 7.5 μm
- center of the cells filled with irregularly shaped, highly refractive mass
- slightly pink to intense pink, sometimes yellowish
- colonies irregularly shaped without sharp outline
- no visible gelatinous sheath
- colonies of about 50 – 250 μm in diameter

No drawings from previous authors available.




Fig. 1: *Rhodobacteria 3*. L = 4 - 7.5 μ m. Part of a slightly squashed colony. All cells are separated from each other. Obj. 100 X.




Fig. 2: *Rhodobacteria 3*. L = 4 – 7.5 μ m. A squashed colony. The cells are embedded in a gelatinous mass without sharp outline. Obj. 100 X.

Rhodobacteria 3
Obj. 100 X




Fig. 3: *Rhodobacteria 3*. L = 4 - 7.5 μ m. A squashed colony. Obj. 100 X.




Fig. 4: *Rhodobacteria 3*. L = 4 – 7.5 μm . A squashed colony. Obj. 100 X.



Fig. 5: *Rhodobacteria 3*. L = 4 – 7.5 μ m. A squashed colony with larger cells. Obj. 100 X..

Rhodobacteria 3
Obj. 100 X



Fig. 6: *Rhodobacteria 3*. L = 4 – 7.5 μm . A slightly squashed colony in brightfield illumination. Obj. 100 X.





Fig. 7 a-b: *Rhodobacteria 3*. L = 4 – 7.5 μ m. The cells in detail. a) In some cells separate spherules are visible instead of the central highly refractive mass. b) In larger cells yellowish droplets are visible in the plasm (arrows) separated from the central highly refractive mass. Obj. 100 X.