

***Songophrya armata* Foissner, 2003**

Most likely ID: n.a.


Synonym: n. a.

Sampling location: [Simmelried](#)

Phylogenetic tree: n.a.

Diagnosis:

- length 90–220 µm, width 60–120 µm
- body ovoid shape, slightly tapered to posterior end
- dorso-ventral slightly flattened
- mouth opening apical, with a ventral mouth slit
- circular oral bulge, mouth slit straight
- row of 5–7 extrusome bundles from anterior to posterior on ventral side
- extrusomes 50–58 µm long, flexible fibers
- on average 102 ciliary rows
- macronucleus ribbon-shaped
- dorsally a field of papillae (5 rows)
- dorsal brush 20 rows (hard to see)
- contractile vacuole terminal
- thickened pellicle with a fringe of 1–2 µm long extrusomes



1 = *Songophrya armata*, after Song and Wilbert

2, 3 = *Pseudoprorodon emmae*, after Kahl

4 = apical view of *Pseudoprorodon emmae*, after Kahl (DB = dorsal brush, DP = dorsal field of papillae)

Songophrya armata

In April 2020 I noticed prostomatid ciliates in an old sample of decomposing plant material from [Simmelried](#). At low magnifications I considered these ciliates to be *Holophrya*. But then I noticed that they were very metabolic and burrowed among the plant material (s. fig. 1 a-c) and that they were dorso-ventrally flattened (s. fig. 2 a-b). These characteristics did not match *Holophrya*. I isolated the 180-220 µm long specimens and examined them more closely at higher magnifications. I immediately noticed a ventral row of bundles of extrusomes extending from the mouth opening to the posterior end (s. figs. 4 and 5). There were between 6-7 of these bundles of extrusomes. The mouth opening was slightly displaced ventrally and surrounded by a circular oral bulge (s. fig. 8). Ventrally, there was also a straight, slit-shaped mouth opening, in the extension of which the bundles of extrusomes were arranged (s. fig. 5). On the dorsal side, immediately behind the mouth opening, there is a 5-row papillae field (s. figs. 6, 7 and 8). I could not clearly identify a dorsal brush. The macronucleus is ribbon-shaped (s. figs. 11 and 12). Probably only one spherical micronucleus is present, but I could not identify it with certainty (s. figs. 10 and 12). The contractile vacuole is located terminally (s. fig. 3a). The extrusomes are 50-58 µm long flexibles fibers (s. figs. 8 and 9). There is a distinctly thickened pellicle visible with a fringe of 1-2 µm long extrusomes (s. fig. 9). I identified the ciliate as *Songophrya armata* based on these characteristics.

Later I realized that I had already photographed this species in August 2008 (also in a sample from [Simmelried](#)), but had failed to recognize it then. After the identification in April 2020 I could find *Songophrya* several times, always among rotting plants from the [Simmrlried](#). I could also notice that the species was especially abundant in old samples that were several weeks old. So far I have detected *Songophrya armata* exclusively in the [Simmelried](#).

The history of the taxonomic classification of *Songophrya armata* is quite complex. When I attempt to assign the ciliate to a known species I came across *Pseudoprorodon emmae* (Berg, 1896) described and drawn by Kahl (s. drawings 2 and 3 above). Kahl also recognized and drew the strange dorsal field of papillae (s. drawing 4 above) of this ciliate and the bundles of extrusomes of the ventral side. However, the species I found has 5 rows of dorsal papillae (s. figs. 6, 7 and 8). Therefore I searched for a redescription of *Pseudoprorodon emmae* and found the description of the Myriokaryonidae by Foissner, 2003 (Foissner, 2003, s. [Literature](#)). In this article Foissner describe *Holophrya* (*Pseudoprorodon*) *emmae* and renamed it to *Berghophrya emmae*. In this article Foissner also shows a drawing of the ventral side with the bundles of extrusomes of *Berghophrya emmae*, which Kahl does not show. The bundles of extrusomes in this species are distributed over a larger field on the ventral side and they are not arranged in a row as in my ciliate. So my ciliate could not be *Berghophrya emmae* (= *Holophrya emmae* = *Pseudoprorodon emmae*) because of this feature and the differently constructed papilla field. However, a further ciliate is also described in the same article. This species was originally described by Song and Wilbert as *Holophrya emmae*. Based on the characteristics described by these authors, Foissner concluded that it is not *Holophrya emmae*, but a new species. This new species, identified solely on the basis of the description by Song and Wilbert, was named *Songophrya armata* by Foissner (Foissner, 2003). Foissner never found *Songophrya armata* himself. Due to this history the species *Songophrya armata* is not currently considered valid. In his article on the Myriokaryonidae, Foissner shows a drawing by Song and Wilber of the ventral side of *Songophrya armata* (s. drawing 1 above). The extrusomal bundles of this species are arranged in a straight longitudinal line, as I have seen in my species. In addition, I could see the circular mouth bulge around the mouth opening (s. figs. 6 and 8), which is also a characteristic of the genus *Songophrya*. Therefore I can confirm these characteristics on the basis of life observations, which were described by Song and Wilbert and which caused Foissner to establish a new species. Since there is also no meaningful alternative, I think the assignment *Songophrya armata* is correct.




Fig. 1 a-c: *Songophrya armata*. L = 220 μ m. A specimen burrows among rotting plants under metabolic deformation. Obj. 40 X.

Songophrya armata
Obj. 40 X



Fig. 2 a-b: *Songophrya armata*. L = 182 μ m. A freely swimming specimen from ventral (a) and lateral from right (b). Obj. 40 X.




Fig. 3 a-b: *Songophrya armata*. L = 182 μ m. Two focal planes of a slightly squashed specimen. On the ventral side (b) the row of extrusome bundles (BE) is visible. CV = contractile vacuole. Obj. 40 X.

Songophrya armata
Obj. 60 X

MS


Fig. 4: *Songophrya armata*. L = 200 μm . A more detailed ventral view of the extrusome bundles running in a straight line from the apical mouth slit (MS) to the anterior end. Obj. 60 X.

Songophrya armata
Obj. 100 X

MS



Fig. 5: *Songophrya armata*. The ventral row of bundles of extrusomes (arrows) in a second specimen. MS = mouth slit. Obj. 100 X.



© Martin Kreutz

Fig. 6: *Songophrya armata*. This semi-apical view shows the arrangement of the dorsal field of papillae (DP) posterior to the mouth opening (MO) and the mouth slit running ventrally and the transition to the row of extrusomal bundles (BE) adjoining it. Obj. 100 X.




Fig. 7: *Songophrya armata*. The dorsal field of papillae (DP) in detail. It consists of 5 rows of 3-6 papillae each. Obj. 100 X.

Songophrya armata
Obj. 60 X




Fig. 8: *Songophrya armata*. A dorsal view on the circular oral bulge (OB) and the dorsal field of papillae (DP). Obj. 100 X.




Fig. 9: *Songophrya armata*. The pharyngeal trichites (PT) in a squashed specimen. There is a fringe of 1-2 μ m long extrusomes (EX) in the thickened pellicle. Obj. 100 X.

Songophrya armata
Obj. 100 X




Fig. 10: *Songophrya armata*. The fiber-shaped, flexible extrusomes with a length of 50 – 58 µm in a strongly squashed specimen. Obj. 100 X.

Songophrya armata
Obj. 60 X



Fig. 11: *Songophrya armata*. A total view of a squashed specimen with the ribbon-shaped macronucleus (Ma). Mi? = probably the spherical micronucleus. Obj. 100 X.




Fig. 12: *Songophrya armata*. The ribbon-shaped macronucleus (Ma) in a second specimen. Mi? = probably the spherical micronucleus. Obj. 100 X.