

## Spathidioides 1

**Most likely ID:** *Spathidioides* nov. spec.

**Synonym:** n.a.

**Sampling location:** [Simmelried](#)

**Phylogenetic tree:** Spathidiida

### Diagnosis:

- body sac-shaped, laterally slightly flattened
- length 110–150 µm
- Oral bulge with dorsal verucciform elevation
- macronucleus elongated ellipsoid
- extrusomes type 1, rod-shaped, 0.40–0.48 µm
- extrusomes type 2, rod-shaped, 0.20–0.24 µm
- about 24–30 longitudinal rows of cilia
- contractile vacuole terminal

No drawings from previous authors available.

In October 2010, I found a specimen of a ciliate among rotting aquatic plants in the [Simmelried](#) with the characteristics of the genus *Spathidioides*. This genus was established by Brodsky (1925) and includes spathidiid ciliates whose oral bulge has on the dorsal side a wart-like elevation. Kahl (1935) described 4 species within this genus:

- *Spathidioides carinata*
- *Spathidioides sulcata*
- *Spathidioides exsecata*
- *Spathidioides armata*

Foissner & Xu (2007) added two more species:

- *Spathidioides euglenivora*
- *Spathidioides rigida*

At the same time, Foissner & Xu note that it is questionable whether separating these species into the genus *Spathidioides* is necessary, as all the aforementioned species have been only very poorly studied. However, since this has not yet happened, I also classify the ciliate I found in this genus.

The ciliate from the [Simmelried](#) is 145 µm long and has a distinctly curved oral bulge, which rises sharply on the dorsal side and ends there in a wart-like protrusion densely equipped with extrusomes (s. fig. 2 a-b). The body is sack- or pouch-shaped and somewhat laterally compressed. The macronucleus is elongated ellipsoid and curved (s. fig. 2 a). It is located in the posterior third. The contractile vacuole is terminal. I was able to clearly identify two types of extrusomes (s. fig. 3). Both are rod-shaped. Type 1 is 4.0–4.8 µm long and has slightly tapered ends, while type 2 is only 2.0–2.4 µm long.


In May 2025, I found a second specimen in the [Simmelried](#) with a length of 118 µm (s. fig. 4 a-c). However, I only found this after placing the coverslip, and due to foreign bodies underneath, I could not further reduce the layer thickness. The habitus, macronucleus, and position of the contractile vacuole, however, correspond to the characteristics of the specimen from 2010.

This combination of features does not match any of the previously described species within the genus *Spathidioides*. Therefore, it must be a previously undescribed, new species, which I provisionally name *Spathidioides* 1.


A similar species was found by Bruce Taylor in April 2022 near Wakefield, Canada (s. [i-Naturalist – Spathidioides](#)). However, he did not provide any information about the size or other characteristics of his find. The similarity to my findings essentially lies in the body shape.




**Fig. 1 a-b:** *Spathidioides* 1. L = 145  $\mu$ m. A freely swimming specimen found in October 2010 in the [Simmelried](#) from right. Obj. 40 X.



**Fig. 2 a-b:** *Spathidioides* 1. L = 145  $\mu$ m. The same specimen as shown in fig. 1 a-b at higher magnification. Note the verruciform elevation (EV) at the dorsal end of the oral bulge (OB). CV = contractile vacuole, Ma = macronucleus. Obj. 60 X.



**Fig. 3:** *Spathidioides* 1. The extrusomes type 1 (EX 1) with a length of 4.2–4.8 µm and the extrusomes type 2 (EX 2) with a length of 2.3–2.4 µm in the strongly squashed specimen as shown in fig. 2 a-b. Obj. 60 X.






**Fig. 4 a-b:** *Spathidioides* 1. L = 118  $\mu$ m. The second specimen found in May 2025 in the [Simmelried](#) from left (a), dorsal (b) and from ventral (c). CV = contractile vacuole, Ma = macronucleus. Obj. 40 X.

In July 2022, I found a third specimen in the [Simmelried](#) (s. fig. 5 a-d). However, this one was significantly smaller at 96  $\mu$ m in length and had 4 spherical macronuclei (s. figs. 7a and 8). The other characteristics, however, matched those of the two specimens shown above. Therefore, this is very likely a post-conjugant, which was in the reorganization phase of the nuclear apparatus. The fact that this is a third specimen of the same species is particularly evidenced by the extrusomes, which are identical in shape and length to those of the specimens shown above (s. fig. 9).



**Fig. 5 a-d:** *Spathidioides* 1. L = 96  $\mu$ m. A third specimen with four macronuclei from left (a, b), dorsal (c) and from ventral (d). The specimen was found in July 2022 in the [Simmelried](#) and is likely a post-conjugant. Obj. 40 X.




**Fig. 6 a-b:** *Spathidioides* 1. L = 96  $\mu$ m. The slightly squashed specimen as shown in fig. 5 a-d from left. Obj. 60 X.




**Fig. 7 a-b:** *Spathidioides 1*. The squashed specimen as shown in fig. 5 a-d from right (a) and from dorsal (b). The four globular macronuclei (Ma) are very likely the result of the reorganisation of the nuclear apparatus after a conjugation. On the dorsal side the dorsal brush (DB) is visible as well as 11 longitudinal rows of cilia (1-11). Obj. 100 X.





**Fig. 8:** *Spathidioides* 1. In the strongly squashed specimen as shown in fig. 5 a-d the four globular macronuclei are visible (Ma 1-4) but no micronucleus. Obj. 100 X.



**Fig. 9:** *Spathidioides* 1. The extrusomes type 1 and type 2 (EX 1, EX 2) in the strongly squashed specimen as shown in fig. 5 a-d. They have the same shape and length as the specimen found in October 2010 (s. fig. 3). Obj. 100 X.