

***Sphaerocystis bavarica***

**(Skuja) Bourrelli, 1966**

**Most likely ID:** n.a.


**Synonym:** *Radiococcus bavaricus*

**Sampling location:** [Simmelried](#)

**Phylogenetic tree:** [Sphaerocystis bavarica](#)

**Diagnosis:**

- spherical or elongated colonies, up to 1 mm
- margin of the colony deliquescent
- young cells almost spherical in tightly arranged groups of 2–4–8 cells
- older cells broadly ovoid or ellipsoidal (6–15 µm long) in widely spaced groups
- cell wall thin and smooth
- during autospore formation, the cell wall disintegrates in colonial mucilage
- single, parietal chloroplast
- one pyrenoid
- asexual reproduction by autospores or zoospores with 2 apical flagella
- benthic lifestyle



after Fott


### *Sphaerocystis bavarica*

*Sphaerocystis bavarica* is one of the most common chlorococcal algae in [Simmelried](#). Especially in pond 6 it occurs in masses since this pond is becoming more and more silted up.


*Sphaerocystis bavarica* forms very large colonies that can contain more than 500 cells and are up to one millimeter in size. The colonies of *Sphaerocystis bavarica* are characterized by the fact that they do not have a sharply defined envelope and that the groups of cells within the colony consist of young, spherical cells as well as older, broadly ovoid cells. With increasing age, the cells move away from each other and the cell wall of the autospores dissolves. This gives the colony a heterogeneous character.

*Sphaerocystis bavarica* can easily be confused with *Sphaerocystis schroeteri* and *Pseudosphaerocystis lacustris*. However, *Sphaerocystis schroeteri* is a planktonic alga with a sharply defined envelope and spherical cells, while *Pseudosphaerocystis lacustris* is a tetrasporal alga with two contractile vacuoles and an eyespot. All these characteristics do not apply to *Sphaerocystis bavarica*. The benthic lifestyle of *Sphaerocystis bavarica* is also an important factor for identification.

*Sphaerocystis bavarica*  
Obj. 20 X




**Fig. 1:** *Sphaerocystis bavarica*. A squashed colony in brightfield illuminations. This colony consists of more than 500 cells. Obj. 20 X.




**Fig. 2:** *Sphaerocystis bavarica*. A slightly squashed second colony. Note that the margin of the colony is not sharply defined (arrows). Obj. 40 X.






**Fig. 3:** *Sphaerocystis bavarica*. A slightly squashed third colony. Note that the remains of the cell walls (RMC). Obj. 40 X.



**Fig. 4:** *Sphaerocystis bavarica*. L = 8–11 µm. Detail of the colony shown in fig. 2. Note that the older cells are not spherical, but broadly oval or ovoid. Obj. 100 X.





**Fig. 5:** *Sphaerocystis bavarica*. L = 5-8 µm. Detail of the colony shown in fig. 1. Note that the younger, smaller cells are almost spherical. PY = pyrenoid. Obj. 100 X.



**Fig. 6:** *Sphaerocystis bavarica*. L = 5–9 µm. Detail of the colony shown in fig. 1 with various stages of autospore formation (FA). SCD = starting cell division. Obj. 100 X.