

Stentor polymorphus
(Müller, 1773) Ehrenberg, 1830

Most likely ID: n.a.


Synonym: n.a.

Sampling location: [Simmelried](#), [Purren pond](#), [Bussenried](#), [Ulmisried](#), [Mühlhalden pond](#)

Phylogenetic tree: [Stentor polymorphus](#)

Diagnosis:

- body elongated trumpet-shaped, contracted specimens ovoid to club-shaped
- length up to 2000 µm (of elongated specimens)
- cytoplasm green due to symbiotic algae (*Chlorella*)
- adoral membranelle running in clockwise direction to oral funnel
- attached to the substrate with thigmotactic cilia
- macronucleus moniliform, 5–20 nodules
- contractile vacuole on left wall of oral funnel



after Kahl

Stentor polymorphus

Stentor polymorphus is a very common ciliate that I find regularly in most of my sampling location. In July 2023, I even observed a mass development of *Stentor polymorphus* in the [Purren pond](#). Billions of specimens covered the top layer of leaves on the mud, turning them an intense green.

Due to its size and green coloration caused by thousands of symbiotic *Chlorella* algae, *Stentor polymorphus* is unmistakable. The macronucleus is monoliform and consists of 5 – 20 nodules. I was able to identify several microsnuclei adjacent to the nodules, which was not previously described by Kahl (1932) or Foissner (1992).

Tactile bristles were observed in *Stentor polymorphus* by Penard (1922). However, this observation has not been confirmed so far. Also I could not recognize any in my

population.



© Martin Kreutz

Fig. 1: *Stentor polymorphus*. L = 550 – 710 μm . A group of four fully extended specimens in darkfield illumination. Obj. 20 X.

Stentor polymorphus
Obj. 40 X



© Martin Kreutz

Fig. 2: *Stentor polymorphus*. L = 510 μ m. A partly extended specimen. Obj. 40 X.




Fig. 3: *Stentor polymorphus*. The adoral zone of membranelles (AZM) in detail. In apical view the adoral zone is almost bretzel-shaped and runs in clockwise direction into the oral funnel (OF). Obj. 40 X.




Fig. 4: *Stentor polymorphus*. Focal plane on the nodules of the moniliform macronucleus (Ma) in a slightly squashed specimen. OF = oral funnel. Obj. 40 X.




Fig. 5: *Stentor polymorphus*. The nodules of the macronucleus (MN) in detail. Note the group of 3 micronuclei (Mi) adjacent to the nodule. SB = symbiotic algae. Obj. 100 X.




Fig. 6: *Stentor polymorphus*. The symbiotic algae have a diameter of 4–6 μm and are member of the genus *Chlorella*. Each *Chlorella* cell has a nucleus (Nu) of its own which is hard to recognize. Obj. 100 X.