

Tropidoatractus acuminatus

Levander, 1894

Most likely ID: n.a.


Synonym: n.a.

Sampling location: [Purren pond, Simmelried](#)

Phylogenetic tree: [*Tropidoatractus acuminatus*](#)

Diagnosis:

- body elongate ellipsoidal, strongly spirally furrowed by 7–8 cortical ridges, shape very variable
- anterior end obliquely truncate, posterior end with distinct spine
- length 70–180 µm
- adoral zone short (11–16 membranelles), right anteriorly
- perizonal stripe longer than adoral zone
- 7–8 rows of cilia running in the furrows of pellicle
- somatic cilia arranged in pairs
- macronucleus elongate ellipsoidal with adjacent, spherical micronucleus
- contractile vacuole at base of spine



1 = after Rottovera et al.

2 = after Kahl

Tropidoatractus acuminatus

I find *Tropidoatractus acuminatus* regularly in the [Purren pond](#) and the [Simmelried](#). The occurrence of this ciliate seems to be quite different. For example, Foissner describes the occurrence as "very rare", while Kahl describes it as "common, never frequent".

Tropidoatractus acuminatus is clearly and easily identifiable by its 7-8 distinct ribs, which run clockwise over the body (s. figs. 1 a and 2 e). The posterior end is elongated to a pointed spine (s. fig. 2 d). The specimens are usually longer than 100 µm. However, *Tropidoatractus acuminatus* is known to be very rich in form. Thus, also strongly broadened, stocky forms can be found and those with a very long terminal spine. In addition, dwarf forms of *Tropidoatractus acuminatus* can be found (see below).

The anterior end is flattened and curved ventrally. On the anterior margin of the apikal dome runs the perizonal stripe, with densely standing, long cilia (s. fig. 2 c). The somatic cilia are arranged in pairs between the ribs (s. fig. 5). In my population the macronucleus was usually quite long and ellipsoidal (s. fig. 4). It was mostly found in the anterior third of the body with an attached, spherical micronucleus (s. fig. 4). I could not detect any symbiotic bacteria in the cytoplasm and none have been described so far. The adoral zone is located on the right side at the anterior end. It is partially covered by the ribs. From the right side, however, it can be seen well (s. fig. 2 a and 2 b).



Fig. 1 a-c: *Tropidoatractus acuminatus*. L = 126 μ m. Three focal planes of a freely swimming specimen from ventral. Obj. 60 X.





Fig. 2 a-e: *Tropidoactractus acuminatus*. L = 142 μ m. A second freely swimming specimen from right (a, b, d) and ventral (c, e). AZM = adoral zone of membranelles, CV = contractile vacuole, Ma = macronucleus, PC = perizonal cilia, PS = posterior spine. Obj. 100 X.

Tropidoatractus acuminatus
Obj. 20 X



© Martin Kreutz

Fig. 3: *Tropidoatractus acuminatus*. Apical view from the posterior end. Note the 8 ribs of this specimen (1-8) running clockwise to the anterior end. Obj. 20 X.

Tropidoatractus acuminatus
Obj. 100 X




Fig. 4: *Tropidoatractus acuminatus*. The macronucleus (Ma) with the adjacent micronucleus (Mi) in a strongly squashed specimen. AZM = adoral zone of

membranelles, RG = accumulation of refractive granules in the apical dome. Obj. 100 X.

Tropidoactractus acuminatus
Obj. 100 X



Fig. 5: *Tropidoatractus acuminatus*. Focal plane on the ventral somatic ciliation of a strongly squashed specimen. The cilia are widely spaced and arranged in pairs (PoC). Obj. 100 X.




Fig. 6: *Tropidoatractus acuminatus*. A specimen during cell division. CV 1, CV 2 = contractile vacuoles, Ma = macronucleus in division. Obj. 100 X.

Tropidoatractus acuminatus can be observed very often during conjugation. Thereby the specimens fuse with the apikal domes and not via the mouth openings. In the area where the specimens are connected an accumulation of refractive granules is visible. It seemed to me that the pellicle of the two specimens also

merged and formed a continuous apical arch on which a continuous perizonal stripe ran (s. fig. 8b).




Fig. 7 a-b: *Tropidoatractus acuminatus*. Two focal planes of a pair in conjugation. Obj. 100 X.



Fig. 8 a-b: *Tropidoatractus acuminatus*. A second pair in conjugation. Obj. 100 X.

Already by Penard and Kahl dwarf forms of *Tropidoatractus acuminatus* were described, in which the terminal spine is reduced. I could also find these forms frequently. Some had a clearly truncated posterior end. The furrows of the pellicle did not run spirally in these forms, but rather straight over the whole body. The length of these forms was often below 50 µm and I found them frequently in conjugation (s. fig. 9 a-b).




Fig. 9 a-b: *Tropidoatractus acuminatus*. A pair of dwarf forms with lengths of 38 μ m and 44 μ m in conjugation. Obj. 100 X.