

Urostomides bacillatus

(Levander, 1894) Bourland et al., 2017

Most likely ID: n.a.


Synonym: *Metopus bacillatus*, *Apometopus bacillatus*

Sampling location: [Purren pond](#), [Bussenried](#), [Simmelried](#)

Phylogenetic tree: [*Urostomides bacillatus*](#)

Diagnosis:

- body obpyriform with prominent apical dome
- length 65–160 µm
- apical dome overhangs left side
- adoral zone arises dorsally, runs almost equatorially to ventral side where it bends posteriorly
- posterior end narrowly rounded or pointed, not tail-like
- macronucleus large, globular with adjacent spherical micronucleus
- prominent fringe of extrusomes (about 3 µm long) beneath pellicle
- contractile vacuole subterminal
- tuft of elongated caudal cilia at posterior end



1 = after Bourland et al.
2, 3 = after Kahl (3 = lateral from left)

Urostomides bacillatus

Urostomides bacillatus was first described in 1894 by Levander as *Metopus bacillatus*. In 2017 the species was redescribed by Bourland et al. who transferred the species to the genus *Urostomides*.

Urostomides bacillatus is very similar to [*Urostomides striatus*](#). However, *Urostomides striatus* is smaller (< 100 µm), has a distinctly tail-like extended posterior end and is also much more dorso-ventrally flattened.

I find *Urostomides bacillatus* regularly and frequently in the mud zone. The species looks somewhat mushroom shaped when free swimming. Close examination is complicated by coverglass sensitivity. However, I was able to determine that a few specimens squeeze easily under the coverslip, while most specimens denature before. The reasons for this different behavior is not known to me.

Kahl mentions that the species is very variable. I can confirm this by the variability of the posterior half of the body. Some specimens were shaped like an ice cream cone, while in other specimens the posterior end appeared distinctly tapered, but this also depended on the degree of filling of the contractile vacuole. This makes it difficult to distinguish *Urostomides bacillatus* from [*Urostomides striatus*](#).

Urostomides bacillatus
Obj. 40 X




Fig. 1 a-f: *Urostomides bacillatus*. L = 100 µm. A freely swimming specimen from dorsal (a, b), ventral (c) and from right (d, e, f). Obj. 40 X.




Fig. 2 a-c: *Urostomides bacillatus*. L = 114 μ m. A second freely swimming specimen from dorsal (a, c) and from left (b). CV = contractile vacuole, MO = mouth opening, UM = undulating membrane. Obj. 60 X.



Fig. 3 a-c: *Urostomides bacillatus*. L = 100 μ m. A third freely swimming specimen from dorsal (a, c) and from right (c). CV = contractile vacuole, FE = fringe of extrusomes, Ma = macronucleus, Mi = micronucleus. Obj. 60 X.

Urostomides bacillatus
Obj. 100 X



© Martin Kreutz

Fig. 4: *Urostomides bacillatus*. L = 124 μ m. A freely swimming specimen from ventral. Obj. 100 X.




Fig. 5 a-b: *Urostomides bacillatus*. L = 128 μ m. A slightly squashed specimen from ventral. AZM = adoral zone of membranelles, MO = mouth opening, UM = undulating membrane. Obj. 100 X.




Fig. 6 a-b: *Urostomides bacillatus*. L = 128 μ m. The same specimen shown in fig. 5 a-b from right (a) and from dorsal (b). Note the distinct striation of the pellicle by the ciliary rows (CR). In the space between the ciliary rows the extrusomes are visible as bright dots. AZM = adoral zone of membranelles, CC = caudal cilia, PC = perizonal cilia. Obj. 100 X.